



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A31K 39/095, C07K 13/00, 7/04 A61K 39/39, 39/385, C12N 15/31 C12N 15/62 // (C12N 15/31 C12R 1/36)	A2	(11) International Publication Number: WO 90/06696 (43) International Publication Date: 28 June 1990 (28.06.90)
(21) International Application Number: PCT/US89/05678 (22) International Filing Date: 19 December 1989 (19.12.89) (30) Priority data: <div style="display: flex; justify-content: space-between;"> <div> 8803111 8900030 8901612 </div> <div> 19 December 1988 (19.12.88) 6 January 1989 (06.01.89) 26 June 1989 (26.06.89) </div> <div> NL NL NL </div> </div> (71) Applicants (for all designated States except US): PRAXIS BIOLOGICS, INC. [US/US]; 30 Corporate Woods, Suite 300, Rochester, NY 14623-1493 (US). RIJKSINSTITUUT VOOR VOLKSGEZONDHEID EN MILIEUHYGINE [NL/NL]; Antonie van Leeuwenhoeklaan 9, P.O. Box 1, NL-3720 BA Bilthoven (NL). (72) Inventors; and (75) Inventors/Applicants (for US only) : SEID, Robert, C., Jr. [US/US]; 590 25th Avenue, San Francisco, CA 94121 (US). PARADISO, Peter, R. [US/US]; 6 Guilford Way, Pittsford, NY 14534 (US). POOLMAN, Jan, T. [NL/NL]; Leeteinde 8, NL-1151 AK Broek in Waterland (NL). HOOGERHOUT, Peter [NL/NL]; Idenburgstraat 13, NL-2805 SZ Gouda (NL). WIERTZ, Emmanuel, J., H., J. [NL/NL]; Mauritsstraat 106, NL-3583 HW Utrecht (NL). VAN DER LEY, Peter [NL/NL]; Adriaan van Ostadelaan 124, NL-3583 AM Utrecht (NL). HECKELS, John, Edward [GB/GB]; 6 Arun Way, West Wellow, Romsey, Hampshire SO51 6GT (GB). CLARKE, Ian, Nicholas [GB/GB]; 15 Fernyhurst Avenue, Rownhams, Southampton, Hampshire SO1 8DR (GB).		(74) Agents: DeCONTI, Giulio, A., Jr. et al.; Hamilton, Brook, Smith & Reynolds, Two Militia Drive, Lexington, MA 02173 (US). (81) Designated States: AT (European patent), AU, BE (European patent), CH (European patent), DE (European patent), DK, ES (European patent), FI, FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), NO, SE (European patent), US. Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: MENINGOCOCCAL CLASS 1 OUTER-MEMBRANE PROTEIN VACCINE (57) Abstract Outer-membrane vesicles, Class 1 outer membrane proteins of <i>Neisseria meningitidis</i> , fragments or oligopeptides containing epitopes of the Class 1 I OMP can be used to immunize against meningococcal disease.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT Austria
AU Australia
BB Barbados
BE Belgium
BF Burkina Fasso
BG Bulgaria
BJ Benin
BR Brazil
CA Canada
CF Central African Republic
CG Congo
CH Switzerland
CM Cameroon
DE Germany, Federal Republic of
DK Denmark

ES Spain
FI Finland
FR France
GA Gabon
GB United Kingdom
HU Hungary
IT Italy
JP Japan
KP Democratic People's Republic
of Korea
KR Republic of Korea
LI Liechtenstein
LK Sri Lanka
LU Luxembourg
MC Monaco

MG Madagascar
ML Mali
MR Mauritania
MW Malawi
NL Netherlands
NO Norway
RO Romania
SD Sudan
SE Sweden
SN Senegal
SU Soviet Union
TD Chad
TG Togo
US United States of America

MENINGOCOCCAL CLASS 1 OUTER-MEMBRANE PROTEIN VACCINE

Description

Background of the Invention

Bacterial meningitis is an inflammatory disease of the central nervous system caused by the growth of bacteria in and adjacent to the leptomeninges. Meningitis is an acute infectious disease which affects children and young adults and is caused by the *Neisseria meningitidis*, amongst other agents including other bacterial and viral pathogens.

Meningococci are subdivided into serological groups depending on the presence of either capsular or cell wall antigens. Currently recognized sero-

groups include A, B, C, D, W-135, X, Y, Z, and 29E as segregated by seroagglutination. The polysaccharides responsible for the serogroup specificity of the group A, B, C, X, W-135 and Y have been

purified.

The carrier rate for meningococci is much higher than the incidence of the disease. Some persons are temporary carriers, while others are chronic carriers, discharging meningococci either more or less continuously or in a sporadic fashion. The meningococcal carrier state is an immunizing process, and within two weeks of colonization, production of antibodies to meningococci can be identified. It appears that bactericidal antibodies are directed against both the capsular polysaccharide and other cell wall antigens.

Studies have shown that meningococcal outer membranes have three to five major proteins, with the predominant 41,000Mr or 38,000Mr proteins carrying the serotype specific determinants. There is a considerable degree of interstrain heterogeneity in the profiles of the outer membrane proteins on sodium dodecyl sulfate-polyacrylamide electrophoretic gels (SDS-PAGE). As defined by peptide mapping studies, the proteins comprise five classes, designated 1 through 5, based upon common peptide structures. Bactericidal monoclonal antibodies have been produced against the 46,000 Mr Class 1 proteins which are shared to some extent among strains of different serotypes. (Frasch, C.E. et al., (1985) pg. 633, "New Developments in Meningococcal Vaccines", in G.K. Schoolnik et al. (ed.) The Pathogenic Nisseriae, American Society for Microbiology, Washington, D.C.).

The capsular polysaccharide of groups A, C, W-135 and Y meningococci have been used to develop vaccines against the organism. Although these vaccines have been effective in the short term, they do not induce immunological memory and subjects must be revaccinated within approximately 3 years to maintain their resistance. The group B polysaccharide is poorly immunogenic and successful vaccines have not been produced. A possible explanation for the low activity may be due to tolerance to the group B polysaccharide induced by crossreactive

antigens found in human tissues such as the brain. Furthermore, studies show that most of the bactericidal antibodies in the convalescent sera of patients who have had group B meningococcal disease are directed against outer membrane proteins.

Vaccines for protecting against group B meningococcal disease have been developed in which non-covalent complexes of outer membrane proteins (OMP) and group B polysaccharide were administered.

Beuvery, et al. (1983) *Infect. Immun.* 40:369-380. However, the B polysaccharide is known to induce a transient IgM antibody response, which does not confer immunoprotection. Furthermore, there is great antigenic diversity and variability in the meningococcal outer membrane proteins from strain to strain. Additionally, lipopolysaccharides are present in the OMP and exhibit antigenic variability as well.

There is a need for safe and effective vaccines against meningococcal disease which provide immunity from infection, particularly in infants and the elderly.

Summary of the Invention

This invention pertains to isolated outer membrane vesicles (OMV's), to substantially purified Class I outer membrane protein (OMP) of *Neisseria meningitidis*, to fragments of the Class I OMP and to oligopeptides derived from the Class I OMP which contain continuous or discontinuous, immunogenic and

protective B cell epitopes reactive with
 bactericidal antibodies against N. meningitidis and
 to the use of isolated OMV's, the meningococcal
 Class I OMP, fragments or oligopeptides for
 vaccination against N. meningitidis.
 The isolated OMV's, meningococcal Class I OMP,
 fragments or oligopeptides derived therefrom can be
 used in univalent or multivalent subunit vaccines
 alone, in mixtures, or as chemical conjugates or
 genetic fusions. In preferred vaccines, epitopes
 from different epidemiologically relevant
 meningococcal strains are used. In addition,
 isolated OMV's, the Class I OMP, fragments or
 oligopeptides can be used in conjugation (as
 mixtures, fusion or conjugates) with other antigens
 of N. meningitidis. For example, they can be used
 in conjugation with capsular polymers or oligomers
 (or fragments thereof) of N. meningitidis or with
 Class I outer membrane proteins (or epitopes thereof)
 of different subtypes. In addition, they can be
 used with antigens of other infectious bacteria,
 viruses, fungi or parasites. Class I OMP cell
 epitopes also are defined and these can be used in
 conjunction with other vaccine components to enhance
 the protective immune response to the vaccines.
 This invention also pertains to the methods of
 producing isolated OMV's, the Class I OMP, fragments
 and oligopeptides and to various vaccine
 formulations containing them. The isolated OMV's
 Class I OMP can be produced by mutant meningococcal
 strains which do not express the Class 2/3

outer membrane protein. Fragments can be produced by cyanogen bromide cleavage and subsequent purification. Isolated OMV's, the class I OMP, fragments or oligopeptides can be produced by recombinant DNA techniques, chemical synthesis or chemical or enzymatic cleavage. These materials, in turn, can be conjugated or fused to carrier peptides or proteins, to other antigens of *N. meningitidis* or to antigens of other microorganisms by chemical or genetic coupling techniques to produce multivalent antigenic conjugates and fusion peptides or proteins. They can be modified for conjugation such as by the addition of amino acids or other coupling groups. For vaccination, isolated OMV's, the class I OMP, fragments or oligopeptides, in any of the forms described, can be formulated in pharmaceutically acceptable vehicles with optional additives such as adjuvants.

This invention also pertains to isolated nucleic acids which encode class I OMP, fragments or oligopeptides. The nucleic acids can be incorporated into appropriate expression systems for production of isolated OMV's, class I OMP, fragments or any oligopeptides derived therefrom. These nucleic acids can be modified as genetic fusions to contain sequences encoding additional polypeptides useful in enhancing the immune response to the vaccine formulation containing the expressed fusion polypeptides. In addition, class I OMP of *N. meningitidis* is homologous in amino acid sequence and structure to porin proteins of other gram negative pathogens and

thus the Class I OMP, fragments and oligopeptides of this invention allow for the development of vaccines for other gram negative pathogens.

Brief Description of the Figures

Figure 1. Scheme for amplification of genes encoding meningococcal Class I outer membrane protein by PCR (Polymerase Chain Reaction).

Figure 2. 5' gene sequences encoding VR1 (first variable region) of Class I outer membrane proteins of several *N. meningitidis* subtypes.

Figure 3. 3' gene sequences encoding VR2 (second variable region) of Class I outer membrane proteins of several *N. meningitidis* subtypes.

Figure 4. Epitope scanning by reaction of monoclonal antibodies with solid phase decapetides spanning the predicted amino acid sequences of Class I proteins from strains P1.7, P1.16 and P1.15. Annotations show the strain from which the sequence was derived, the map used and its subtype specificity.

Figure 5. Reaction of the monoclonal antibodies with series of overlapping decapetides corresponding to variable regions VR1 and VR2, with adjacent peptides differing by a single amino acid residue. Annotations show the strain from which the sequence was derived, the map used and its subtype specificity.

This invention also comprises the application of genetic engineering with the object of producing isolated OMV's Class I OMP or portions thereof, cloned DNA in recombinant DNA expression vectors. This invention also comprises the application of pure Class I outer-membrane proteins with the aid of membrane protein; the preparation of isolated OMV's strains which do not express the Class 2/3 outer-Class I outer-membrane proteins, using mutant of the OMP; the preparation of isolated OMV's, pure cyanogen bromide) and oligopeptides bearing epitopes of the OMP (e.g., prepared by the application of isolated OMV's, meningococcal Class I OMP, fragments This invention pertains to vaccines comprising

Detailed Description of the Invention

Meningitidis Class I OMP subtype Pl.16. Figure 11. Putative conformation of N. Meningitidis Class I OMP conjugated to CRM 197. analyses of a conjugate comprising an epitope of N. Figure 10. Representative Western blot of recombinant flagellin. Figure 9. Representative analysis by SDS-PAGE of recombinant flagellin. Figure 8. Representative chromatogram of high performance liquid chromatography of a recombinant Meningitidis Class I OMP subtype Pl.6,16. expressing variable region epitopes of N. Figure 7. Structure of recombinant flagellins Meningitidis Class I OMP subtype Pl.6,16. flagellins expressing variable region epitopes of N. Figure 6. Construction of recombinant

genetic fusions of Class I OMP, portions or epitopes thereof; and the preparation of multivalent Class I outer-membrane vaccine through peptide synthesis, as the epitopes with a short peptide chain can be synthetically prepared.

It has emerged that meningococcal Class I outer-membrane proteins induce a strong bactericidal immune response to the strains containing the appropriate subtype epitopes, irrespective of whether these are from group A, B, C, W-135, and Y strains. The polysaccharide vaccine can be enhanced or replaced by a vaccine according to the invention as a vaccine with broad, extensive action against most serotypes. The protective bactericidal monoclonal antibodies specific for the Class I outer-membrane protein react strongly with fragments that have been split off and short synthetic peptides which have been prepared using the amino acid sequence of Class I outer-membrane proteins. Since meningococcal disease is currently caused chiefly by Group B meningococci and because the Class I outer-membrane proteins occurring in Group B meningococci also occur in group A, C, W-135, Y meningococci, vaccines of this invention which comprise one or more Class I OMP epitopes derived from N. meningitidis Group B should be effective in preventing disease caused by group A, C, W-135 and Y. Preferably, the preparation of such a vaccine starts from at least two different immunogenic and protective epitopes which have been selected on

epidemiological grounds. Vaccines according to the invention comprise, for example, at least one protein which is obtained either in OMV formulation or by purification from mutant strains producing one or more class I OMP or at least two fragments prepared through a cyanogen bromide fragmentation or at least two synthetic peptides, chosen from about 10 major epitopes, or products obtained by gene expression via recombinant DNA technology, which contain the desired epitopes. To maximize efficacy to a broad range of meningococcal strains, the greater number of different protective epitopes in the vaccine the better. In addition, the vaccines according to the invention may advantageously contain meningococci A and C or optionally W-135 and Y polysaccharides and/or detergents. Preferably, the A and C polysaccharides are covalently coupled to a protein or polypeptide carrier. These carriers include, for example, isolated OMV, the class I OMP protein, T-helper epitopes, bacterial toxins, toxoids, nontoxic mutants (CRM's), recombinant *Salmonella* flagellin and viral particles such as rotavirus VP6 protein, Hepatitis B surface antigen or parvovirus VP1 and VP2 proteins. Both Zwitter-ionic, cationogenic, anionogenic and nonionic detergents are Zwittergent 3-10, Zwittergent 3-14 (N-tetradecyl-N, N-dimethyl-3-aminomethyl-propane sulfonate), Tween-20, sodium deoxycholate, sodium cholate and octylglucoside. The vaccines according to the invention may also contain an adsorbent such

as aluminum hydroxide, calcium phosphate; or advantageously, aluminum phosphate. The fragments, proteins, peptides can also be processed in immunostimulating complexes (ISCOMS), liposomes or microspheres for delivering and/or use as an adjuvant or in connection with other adjuvants so that greater immunogenicity is obtained.

This invention encompasses isolated OMV, substantially pure meningococcal class 1 outer membrane proteins (of any subtype) and fragments of the proteins containing epitopes thereof. The fragments can be any portions of the molecular weight of 25kd or less which contain epitopes which are bound by protective bactericidal antibodies against *N. meningitidis*. These include proteolytic fragments and synthetic oligopeptides which are comprised of amino acid sequences which correspond, at least in part, to epitopes of a Class I OMV. The isolated OMV's, Class I OMV, fragments or epitope-containing oligopeptides derived therefrom can be comprised of amino acid sequences which are different, but essentially biologically equivalent to the natural sequences. These sequences can include sequences in which functionally equivalent amino acid residues are substituted for residues within the sequence resulting in a silent change. For example, one or more amino acid residues within the sequence can be substituted by another amino acid of a similar polarity which acts as a functional equivalent, resulting in a silent alteration.

Substituted amino acids within the sequence may be selected from other members of the class to which the amino acid belongs. For example, the nonpolar (hydrophobic) amino acids include glycine, alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan and methionine. The polar neutral amino acids include serine, threonine, cysteine, tyrosine, asparagine, and glutamine. The charged (basic) amino acids include arginine, lysine and histidine. The negatively charged (acidic) amino acids include aspartic and glutamic acids. Additionally, isolated OMV's, the class I OMP, fragments or the oligopeptides can be modified for conjugation to other molecules (e.g., by the attachment of coupling groups such as the amino acids cysteine and/or lysine or other linking groups and/or spacer groups) including other class I OMP of a different subtype, T cell epitopes, B cell epitopes, carrier peptides or proteins or adjuvanting molecules. As described in detail below, the class I OMP. fragments or oligopeptides can be used in many different forms (e.g., alone, in mixtures, or as conjugates and genetic fusions produced from recombinant DNA vectors) in vaccines. For these purposes, the materials can be produced by isolation from *N. meningitidis*, by proteolytic digestion, by chemical synthesis, or by expression as recombinant molecules. The methods of production and use of the isolated OMV's, the class I OMP and the fragments and the oligopeptides of class I OMP are described below.

Protein modeling and structure analysis of the Class I OMPs were performed using the principles for several *E. coli* outer membrane proteins. (Vogel, H. et al., J. Mol. Bio., 190:191 (1986); Ference, T. et al., J. Mol. Bio., 201:493 (1988) and Tomassen, J. in "Membrane Biogenesis", NATO ASI Series H16, pp351, Springer-Verlag, NY (1988)). The derived amino acid sequence of the Class I OMPs were used for the modeling studies and comparison. The amino acid sequence homology was compared to other gram negative bacterial porin proteins and similarity was established for the protein structure. Exposed surface loops and transmembrane structure were very similar for these porin proteins. With the information revealed concerning variable and constant region protective epitopes of N. meningitidis and their structure, one can predict based upon the amino acid sequence where protective epitopes may reside for other pathogenic gram negative bacteria to be evaluated and included in vaccines for the same.

Production of isolated OMV's

OMV's can be produced either from the culture supernatant or from the bacterial cells after fragmentation as described by Beuvery et al. (1983) loc. cit. OMV's carrying proteins from more than one meningococcus can be isolated from strains manipulated to express heterologous proteins.

-13-

Production and Purification of Class I OMP and CNBr fragments thereof

Class I and Class 3 outer membrane proteins can be isolated as described by Beuvery, E.C. et al., Antonie van Leeuwenhoek J. Microbiol. 52:232 (1986). The production of substantially pure Class I OMP free of Class 2 or 3 OMP's is achieved by this method using mutant meningococcal strains which do not express Class 2/3 OMP. A preferred strain for production of Class I OMP is the H115 strain, deposited as CBS 636.89.

Fragments can be produced by cyanogen bromide cleavage as described by Teerlink T. et al., J. Exp. Med. 166:63 (1987) for a gonococcal protein. The N-terminal fragment is referred to as CB-1 and the C-terminal fragment is referred to as CB-2. These CNBr fragments can be purified via reverse phase HPLC on a VydaxTM C4 or an AquaporTM R-300 column using a water/acetonitrile gradient. Alternatively, the fragment can be purified by multiple cold trichloroacetic acid precipitations. These procedures remove greater than 95% of interfering contaminants (e.g., buffer salts, detergents and fragment contaminants).

Preparation of fragments and oligopeptides containing epitopes of class I OMP

A. Preparation by proteolytic digestion
Oligopeptides containing epitopes reactive with bactericidal antibodies against *N. meningitidis* can be produced by digestion of the class I OMP, CB-1 or CB-2 fragments with proteinases such as endolys-C,

-14-

endocytosis, endocytosis and staphylococcus V8-protease. The digested fragments can be purified by, for example, high performance liquid chromatographic (HPLC) techniques.

B. Preparation by chemical synthesis

Oligopeptides of this invention can be synthesized by standard solid peptide synthesis (Barany, G. and Merrifield, R.B., The Peptides 2:1-284, Gross, E. and Meienhofer, J., Eds., Academic Press, New York) using tert-butyloxycarbonyl amino acids and phenylacetamidomethyl resins (Mitchell, A. R. et al., J. Org. Chem. 43:2845-2852 (1978)) or 9-fluorenylmethoxycarbonyl amino acids on a polyamide support (Dryland, A. and Sheppard, R.C., J. Chem. Soc. Perkin Trans. I, 125-137 (1986)). Alternatively, synthetic peptides can be prepared by pepsin synthesis (Geysen, H.M. et al., J. Immunol. Methods 03:259 (1987); Proc. Natl. Acad. Sci. USA 81:3998 (1984)), Cambridge Research Biochemicals, Cambridge, U.K. or by standard liquid phase peptide synthesis. The deletion or substitution of amino acids (and including extensions and additions to amino acids) in other ways which do not substantially detract from the immunological properties of the oligopeptide.

C. Preparation by recombinant DNA techniques

The Class I OMP, fragments and oligopeptides which exhibit epitopes of the Class I OMP can be

produced by recombinant DNA techniques. In general, these entail obtaining DNA sequences which encode the desired OMP, [Barlow et al., (1989) Mol. Micro., 3:131] fragment or oligopeptide sequences and introducing into an appropriate vector/host expression system one or more similar or different DNA sequences of class I OMP's where it is expressed. The DNA can consist of the gene encoding the class I OMP or any segment of the gene which encodes a function- al epitope of the OMP. The DNA can be fused to DNA encoding other antigens of *N. meningitidis* (such as other outer membrane proteins either of the same or different class) or antigens of other bacteria, viruses, parasites or fungi to create genetically fused (sharing a common peptide backbone), multi- valent antigens. For example, class I OMP fragments can be fused to another class I outer membrane protein of a different subtype (or fragments or epitopes thereof) of *N. meningitidis* to yield fusion proteins comprising multiple class I outer membrane protein subtype determinants.

Genetic engineering techniques can also be used to characterize, modify and/or adapt the encoded peptides or proteins. For example, site directed mutagenesis to modify an OMP fragment in regions outside the protective domains, for example, to increase the solubility of the subfragment to allow easier purification. DNA can also be manipulated to effect superproduction of OMP fragments or combinations thereof in various organisms.

the promoter and initiates transcription of a gene expression vector. RNA polymerase normally binds to cloned DNA, a promoter must be present in the In order to obtain efficient expression of the elements can be used.

number of suitable transcription and translation upon the host-vector system utilized, any one of a vary in their strength and specificities. Depending virus). The expression elements of these vectors cell systems infected with virus (e.g., baculo- (e.g., vaccinia virus, adenovirus, etc.); insect vectors; mammalian cell systems infected with virus DNA; microorganisms such as yeast containing yeast formed with bacteriophage DNA, plasmid DNA or cosmid are not limited to the following: bacteria trans- the host cell used. Host-vector systems include but Primarily the vector system must be compatible with express the oligopeptides of this invention. A variety of host-vector systems can be used to described and the amino acid sequences deduced.

different subtypes can be obtained by the procedures herein. Related DNA sequences for class I OMP of 155:335-350, using the primer sequences disclosed Mullis and Faloona, (1987) Method. Enzym. amplified from bacterial DNA by the methods of Mol. Micro. 3:131 (1989). Class I OMP genes can be Immune 55:2734-40 (1987) and Barlow, A.K. et al., ed as described by Barlow, A.K. et al. Infect. peptides can be synthesized or isolated and sequenc- DNA nc ding a Class I OMP, fragments or oligo-

or a group of linked genes and regulatory elements (called an operon). Promoters vary in their "strength", i.e., their ability to promote transcription. It is desirable to use strong promoters in order to obtain a high level of transcription and, hence, a high level of DNA expression. Depending upon the host cell system any one of a number of suitable promoters can be used. For instance, for *E. coli*, its bacteriophages or plasmids, promoters such as the λ g promoter, λ PR promoter, λ gA promoter, ribosomal RNA promoter, and P_L or P_R promoters of coliphage lambda and others including but not limited to λ gUV5, λ gMPF, λ gA, λ gP and the like, may be used to direct high levels of transcription of adjacent DNA segments. Additionally, a hybrid λ PR- λ gUV5 (tac) promoter or other *E. coli* promoters produced by recombinant DNA or other synthetic DNA techniques may be used to provide for transcription of the inserted DNA.

Bacterial host cells and expression vectors may be chosen which inhibit the action of the promoter unless specifically induced. In certain operons the addition of specific inducers is necessary for efficient transcription of the inserted DNA; for example, the λ g operon is induced by the addition of lactose or IPTG (isopropylthio-beta-D-galactoside). A variety of other operons, such as λ PR, etc., are under different controls. The λ PR operon is induced when tryptophan is absent in the growth media; and the P_L promoter of lambda can be induced by an

increase in temperature in host cells containing a temperature sensitive lambda repressor, e.g., c1857. In this way, greater than 95% of the promoter-directed transcription may be inhibited in unduced cells. Thus, expression of the recombinant peptide or protein can be controlled. This is important if the expression product of the DNA is lethal or detrimental to the host cells. In such cases, transformants may be cultured under conditions such that the promoter is not induced; then, when the cells reach a suitable density in the growth medium, the promoter can be induced for production of the protein.

One such promoter/operator system is the "tac" or trp-lac promoter/operator system (Russell and Bennett, 1982, Gene 20:2312-243; DeBoer, European Patent Application, 67, 540 filed May 18, 1982). This hybrid promoter is constructed by combining the -35 b.p. (-35 region) of the trp promoter and the -10 b.p. (-10 region or Pribnow box) of the lac promoter (the sequences of DNA which are the RNA polymerase binding site). In addition to maintaining the strong promoter characteristics of the tryptophan promoter, lac is also controlled by the lac repressor.

When cloning in a eukaryotic host cell, enhancer sequences (e.g., the 72 bp tandem repeat of SV40 DNA or the retroviral long terminal repeats or LTRs, etc.) may be inserted to increase transcriptional efficiency. Enhancer sequences are a set of

eucaryotic DNA elements that appear to increase transcriptional efficiency in a manner relatively independent of their position and orientation with respect to a nearby gene. Unlike the classic promoter elements (e.g., the polymerase binding site and the Goldberg-Hogness "TATA" box) which must be located immediately 5' to the gene, enhancer sequences have a remarkable ability to function upstream from, within, or downstream from eucaryotic genes; therefore, the position of the enhancer sequence with respect to the inserted DNA is less critical.

Specific initiation signals are also required for efficient gene transcription and translation in procaryotic cells. These transcription and translation initiation signals may vary in "strength" as measured by the quantity of gene specific messenger RNA and protein synthesized, respectively. The DNA expression vector, which contains a promoter, may also contain any combination of various "strong" transcription and/or translation initiation signals. For instance, efficient translation in *E. coli* requires a Shine-Dalgarno (SD) sequence about 7-9 bases 5' to the initiation codon (ATG) to provide a ribosome binding site. Thus, any SD-ATG combination that can be utilized by host cell ribosomes may be employed. Such combinations include but are not limited to the SD-ATG combination from the *cis* gene or the *N* gene of coliphage lambda, or from the *E. coli* tryptophan E, D, C, B or A genes.

-20-

Additionally, any SD-ATC combination produced by

recombinant DNA or other techniques involving

incorporation of synthetic nucleotides may be used.

Any of the methods described for the insertion

of DNA into an expression vector can be used to

ligate a promoter and other genetic control elements

into specific sites within the vector. *N. meningi-*

tidis sequences for expression can be ligated into

an expression vector at a specific site in relation

to the vector promoter and control elements so that

when the recombinant DNA molecule is introduced into

a host cell the foreign genetic sequence can be

expressed (i.e., transcribed and translated) by the

host cell.

The recombinant DNA vector can be introduced

into appropriate host cells (bacteria, virus, yeast,

mammalian cells or the like) by transformation,

transduction or transfection (depending upon the

vector/host cell system). Host cells containing the

vector are selected based upon the expression of one

or more appropriate gene markers normally present in

the vector, such as ampicillin resistance or tetra-

cycline resistance in pBR322, or thymidine kinase

activity in eucaryotic host systems. Expression

vectors may be derived from cloning vectors, which

usually contain a marker function. Such cloning

vectors may include, but are not limited to the

following: SV40 and adenovirus, vaccinia virus

vectors, insect viruses such as baculoviruses, yeast

vector, bacteriophage vectors such as lambda gt-WES-

- 21 -

Lambda B, Charon 28, Charon 4A, Lambda gt-1-lambda BC, Lambda gt-1-lambda B, M13mp7, M13mp8, M13mp9, or plasmid DNA vectors such as pBR322, pAC105, pVA51, pACYC177, pKH47, pACYC184, pUB110, pMB9, pBR325, Col E1, pSC101, pBR313, pML21, RSF2124, pCR1, RP4, pBR328 and the like.

Expression vectors containing the DNA inserts can be identified by three general approaches: (1) DNA-DNA hybridization using probes comprising sequences that are homologous to the inserted gene; (2) presence or absence of "marker" gene functions (e.g., resistance to antibiotics, transformation phenotype, thymidine kinase activity, etc.); and (3) expression of inserted sequences based on the physical immunological or functional properties of the gene product.

Once a putative recombinant clone which expresses a desired class I OMP amino acid sequence is identified, the gene product can be analyzed as follows. Immunological analysis is especially important because the ultimate goal is to use the gene products in vaccine formulations and/or as antigens in diagnostic immunoassays. The expressed peptide or protein should be immunoreactive with bactericidal antibodies against *N. meningitidis*. This reactivity may be demonstrated by standard immunological techniques, such as radioimmuno-precipitation, radioimmune competition, ELISA or immunooblots.

Once the gene product is identified as a class I OMP fragment or an oligopeptide contain a functional epitope thereof, it can be isolated and purified by standard methods including chromatography (e.g., ion exchange, affinity, and sizing column chromatography), centrifugation, differential solubility, or by any other standard techniques for the purification of proteins. Several techniques exist for purification of heterologous protein from prokaryotic cells. See e.g., Olson, U.S. Patent No. 4,518,526, Wetzel, U.S. Patent No. 4,599,197 and Hung et al., U.S. Patent No. 4,734,362. The purified preparation however produced should be substantially free of host toxins which might be harmful to humans. In particular, when expressed in gram negative bacterial host cells such as *E. coli* or *Salmonella*, the purified peptide or protein should be substantially free of endotoxin contamination.

Class I OMP, fragments and oligopeptides of this invention can be formulated as univalent and multivalent vaccines. These materials can be used as produced or isolated by the methods described above. They can be mixed, conjugated or fused with other antigens, including B or T cell epitopes of other antigens. In addition, they can be conjugated to a carrier protein as described below for oligopeptides.

When a haptenic oligopeptide is used (i.e., a peptide which reacts with cognate antibodies, but

cannot itself elicit an immune response), it can be conjugated to an immunogenic carrier molecule. Conjugation to an immunogenic carrier can render the oligopeptide immunogenic. The conjugation can be performed by standard procedures. Preferred carrier proteins for the haptenic oligopeptides are toxins, toxoids or any mutant crossreactive material (CRM) of the toxin from tetanus, diphtheria, pertussis, *Pseudomonas*, *E. coli*, *Staphylococcus*, and *Streptococcus*. A particularly preferred carrier is CRM¹⁹⁷ of diphtheria toxin, derived from *P. diphtheriae* strain C7(β 197) which produces CRM¹⁹⁷ protein. This strain has ATCC accession no. 33281. Alternatively, a fragment or epitope of the carrier protein or other immunogenic protein can be used. For example, the hapten can be coupled to a T cell epitope of a bacterial toxin, toxoid or CRM. See U.S. Patent Application Serial No. 150,688, filed February 1, 1988, entitled "Synthetic Peptides Representing a T-Cell Epitope as a Carrier Molecule For Conjugate Vaccines", the teachings of which are incorporated herein. Other carriers include viral particles composed of Rotavirus VP6, Hepatitis B surface antigen or parvovirus VP1 and VP2. The peptides or proteins of this invention can be administered as multivalent subunit vaccines in combination with antigens of *N. meningitidis* or antigens of other organisms. Some of the other organisms include the pathogenic bacteria *H. influenzae*, *N. meningitidis*, *B. catarrhalis*, *N.*

Bonorrhoeae, *E. coli*, *S. pneumoniae*, etc. For example, they may be administered in conjunction with oligo- or polysaccharide capsular components of *N. meningitidis*. The capsular components can be derived from any of the serological groups, including A, B, C, D, X, Y, Z, 29E and W135. Class I outer membrane proteins of different subtypes can be used. These may be used in combination to evoke bactericidal antibodies against *N. meningitidis*. For example, a fragment derived from class I outer membrane protein of the P1.7.16 subtype can be used together with outer membrane proteins or fragments of outer membrane proteins of other subtypes, such as P1.1, P1.1.16; P1.2; P1.6; P1.9; P1.15; P1.16; or P1.4 (Abdillahi, H. et al. 1988 *Micro. Pathog.* 4:27) or with meningococcal polysaccharides in mixtures or as chemical conjugates. For combined administration with epitopes of other outer membrane proteins, they can be administered separately, as a mixture or as a conjugate or genetic fusion peptide or protein. The conjugates can be formed by standard techniques for coupling proteinaceous materials or techniques for coupling saccharide polymers to proteins. Fusions can be expressed from fused gene constructs prepared by recombinant DNA techniques as described. As mentioned, class I OMP, fragment or any oligopeptides derived therefrom can be used in conjunction with antigens (e.g., polymer capsules or saccharide units, envelope or surface proteins) of

other pathogenic organisms (e.g. bacteria (encapsulated or nonencapsulated), viruses, fungi and parasites). Additional examples of other organisms include respiratory syncytial virus, rotavirus, malaria parasites, and *Cryptococcus neoformans*. In formulating the vaccine compositions with the peptide or protein, alone or in the various combinations described, the immunogen is adjusted to an appropriate concentration and formulated with any suitable vaccine adjuvant. Suitable adjuvants include, but are not limited to: surface active substances, e.g., hexadecylamine, octadecylamine, octadecyl amino acid esters, lysolactin, dimethyl-dioctadecylammonium bromide), methoxyhexadecylglycerol, and pluronic polyols; polyamines, e.g., pyran, dextran sulfate, poly IC, carbopol; peptides, e.g., muramyl dipeptide and derivatives, dimethylglycine, cutisin; oil emulsions; and mineral gels, e.g., aluminum hydroxide, aluminum phosphate, etc., lymphokines and immune stimulating complexes (ISCOMS). The immunogen may also be incorporated into liposomes, microspheres, or conjugated to polysaccharides and/or other polymers for use in a vaccine formulation.

The vaccines can be administered to a human or animal in a variety of ways. These include intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, oral and intranasal routes of administration.

Live vaccines

The peptide and proteins of this invention can be administered as live vaccines. To this end, recombinant microorganisms are prepared that express the peptides or proteins. The vaccine recipient is inoculated with the recombinant microorganism which multiplies in the recipient, expresses the Class I OMP, fragment or oligopeptide thereof and evokes a immune response to *N. meningitidis*. Live vaccine vectors include: adenovirus, cytomegalovirus and preferably, pox viruses such as vaccinia (Paolletti and Panicali, U.S. Patent No. 4,603,112) and attenuated *Salmonella* strains (Stocker, U.S. Patent No. 4,550,081 and Curtiss et al., Vaccine 6:155-160 (1988)). In addition, class I OMP epitopes can be incorporated into the flagella of attenuated bacterial strains.

Live vaccines are particularly advantageous because they lead to a prolonged stimulus which can confer substantially long-lasting immunity. When the immune response is protective against subsequent *N. meningitidis* infection, the live vaccine itself may be used in a preventative vaccine against *N. meningitidis*.

Multivalent live vaccines can be prepared from a single or a few recombinant microorganisms that express different epitopes of *N. meningitidis* (e.g., other outer membrane proteins from other subtypes or epitopes thereof). In addition, epitopes of other

pathogenic microorganisms can be incorporated into the vaccine. For example, a vaccinia virus can be engineered to contain coding sequences for other epitopes in addition to those of *N. meningitidis*. Such a recombinant virus itself can be used as the immunogen in a multivalent vaccine. Alternatively, a mixture of vaccinia or other viruses, each expressing a different gene encoding for different epitopes of outer membrane proteins of *N. meningitidis* and/or epitopes of other disease causing organisms can be formulated as a multivalent vaccine. An inactivated virus vaccine may be prepared. Inactivated vaccines are "killed", i.e., infectivity has been destroyed, usually by chemical treatment (e.g., formaldehyde treatment). Ideally, the infectivity of the virus is destroyed without affecting the proteins which carry the immunogenicity of the virus. In order to prepare inactivated vaccines, large quantities of the recombinant virus expressing the desired epitopes are grown in culture to provide the necessary quantity of relevant antigens. A mixture of inactivated viruses expressing different epitopes may be used for the formulation of "multivalent" vaccines. In certain instances, these "multivalent" inactivated vaccines may be preferable to live vaccine formulation because of potential difficulties arising from mutual interference of live viruses administered together. In either case, the inactivated virus or mixture of viruses may be formulated in a suitable adjuvant in

Type specific monoclonal antibodies were prepared against various meningococci Class I outer-membrane proteins. These monoclonal antibodies recognize the following subtypes: P1,1; P1,2; P1,6; P1,7; P1,9; P1,10; P1,15; P1,16; and P1,17 (now called P1,14). The monoclonal antibodies are available as "Monoclonal Kit Serotyping Meningococci" from the RIVM, Bilthoven, The Netherlands. All these monoclonal antibodies react with the SDS (sodium dodecyl sulphate) denatured protein when tested by Western blotting. It also emerged that a number of these monoclonal antibodies reacted with a

EXAMPLE 1: Monoclonal Antibodies Against Class I OMP's and Their Biological Activity

EXEMPLIFICATION

order to enhance the immunological response to the antigens. Suitable adjuvants include: surface active substances, e.g., hexadecylamine, octadecylamine acid esters, octadecylamine, lysolactin, dimethyl-dioctadecylammonium bromide, N, N-dioctadecyl-N', N'-bis (2-hydroxyethyl-propane diamine), decyl-N', N'-bis (2-hydroxyethyl-propane diamine), methoxyhexadecylglycerol, and pluronic polyols; polyamines, e.g., pyran, dextran sulfate, poly IC, carbopol; peptides, e.g., muramyl dipeptide and derivatives thereof, dimethylglycine, tufsin; oil emulsions; and mineral gels, e.g., aluminium hydroxide, aluminium phosphate, and lymphokines.

25Kd CNBr fragment of the 42Kd Class 1 outer-membrane protein (see below). This result implied that the Class 1 outer-membrane protein epitopes are mainly of linear type and can therefore be copied with synthetic peptides. The epidemiological results of tests carried out by the Applicants show that the described monoclonal antibodies can subtype most of the group A,B,C meningococci which suggests a limited heterogeneity. Each Class 1 outer membrane protein also appears to contain two individual type specific epitopes (Abdillahi and Poolman, Microb. Pathogen, 1988, 4: pages 27-32; idem FEMS Microbiol. Immunol. 47: pages 139-144).

The purified Class 1 outer-membrane protein (see below), subtype P1,7.16, originating from the culture of the Class 2/3 free mutant (H1115) appeared to induce a bactericidal antibody response of 1:64 serum dilution in a dose of 2.5 µg in mice. The monoclonal antibodies against meningococci Class 1 outer-membrane proteins, Class 2/3 outer-membrane proteins and lipopolysaccharides were compared as to bactericidal effect. The monoclonal antibodies against the Class 1 outer-membrane proteins appeared to possess the strongest bactericidal activity (see Table 1). The bacterial response was determined as per Poolman, J.T. (1985), in Schoonick, G.J. et al. Eds. 'The Pathogenic Neisseriae' ASM Publications, Washington, D.C., page 562.

TABLE 1

Bactericidal activity of a collection of monoclonal antibodies, directed against the class 1 (cl 1), class 2/3 (cl 2/3) and lipopolysaccharide (LPS) of meningococci. (ND = not determined).

Test strain	Bactericidal activity of antibody pool (titre)		
strain (Gp:serotype: subtype:LPS type)	cl 2/3 pool	cl 1 pool	LPS pool

3006 (B:26:Pl.2:L2)	1000	8000	ND
M981 (B:4Pl.-:L5)	10	ND	2000
M990 (B:6:Pl.6:L/)	10	2000	ND
M978 (B:8:Pl.1:L1.8)	ND	8000	1000
M982 (B:9:Pl.9:L3.7)	500	2000	1000
M355 (B:15:Pl.15:L1.8)	1000	8000	1000
H44/76 (B:15:Pl.7.16:L3.7)	1000	8000	4000

-31-

The bactericidal activity of these monoclonal antibodies appears to correlate well with the *in vivo* protective activity as measured in the rat meningitis model of Saukkonen et al., 1987, Microbial Pathogen 3:261.

EXAMPLE 1A: Construction of meningococcal strains carrying multiple Class I genes

Replacement of chromosomal genes by clones, slightly different versions has been described for *Neisseria gonorrhoea*. (Stein, D.C., Clin. Microbiol. Rev. 2 (Suppl.), S146-S149 (1989).) We have found that this method can be applied to the Class I gene in *Helicobacter meningitidis*. This was done in the following way:

(1) The Class I gene of strain 2996 (subtype PI.2) was cloned into the vector pTZ19R. (Mead, D.A. et al., Protein Engineering 1, 67 (1986).) The complete gene is located on a 2.2 kb XbaI fragment that was ligated to XbaI digested vector DNA.

(ii) The resulting plasmid was used for transformation of strain H44/76 (subtype PI.7.16). Cells of the acceptor strain were incubated with plasmid DNA in the presence of Mg^{2+} and normal meningococcal medium; they were subsequently diluted and plated, and the

resulting colonies were tested for their ability to bind PL.2-specific monoclonal antibody. Such transformants were found with a frequency of approximately 10^{-3} . Further characterization showed that replacement of the H44/76 Class 1 gene had indeed occurred. An essential feature of the method is the presence of the donor gene on a circular plasmid DNA molecule that is not able to replicate in *N. meningitidis*, since the use of linearized DNA yielded no transformants at all. Construction of a strain with two Class 1 genes was done by a modification of the method described above. For this purpose, the PL.2 Class 1 gene was inserted into a clones Class 5 gene. The Class 5 gene family has two features which make it particularly suitable for this construction.

(Meyer, T.F. and Van Putten, J.P.M., *Clin. Microbiol. Rev.*, 2 (Suppl.) S139-S145 (1989): (1) there are four or five Class 5 genes present in the meningococcal genome, and (ii) expression of these genes is not necessary for growth under laboratory conditions. A Class 5 gene was cloned from strain H44/76 and the PL.2 gene was inserted into an *SphI* site located in or very close to the Class 5 gene. The resulting hybrid plasmid, pMC22, was used for transformation of strain H1115, a Class 3-deficient mutant of H44.76. Colonies reacting with the PL.2-specific monoclonal antibody were isolated and

The purification is carried out according to Beuvery *et al.* (1983) loc. cit. This culture can be done with the desired wild type strains, mutant meningococci strains without Class 2/3 outer-membrane proteins and/or homologous

EXAMPLE 1B: Purification of isolated OMV's from bacteriological culture

purified Class I OMVs. strains can be used to prepare mixtures of different be clones and used separately. These recombinant transformation step, the different Class 5 genes can Class 5 gene can be used in each subsequent with four or five different Class I genes. The same Class I subtypes, it is possible to make a strain By continuing this construction with other the acquisition of a second Class I gene. protein and by Southern blotting, which demonstrated which revealed the presence of both types of Class I sequences on plasmid and chromosome must have that recombination between the Class 5 gene Class I subtypes, i.e., P1.7,16 and P1.2, suggesting However, one transformant was found which made both Class I genes, resulting in subtype replacement. cases recombination has only occurred between the acceptor strain. This indicates that in all these were found to have lost the P1.16 epitope of the characterized. Out of 10 such transformants, nine

and heterogeneous recombinant microorganisms which

express one or more of the desired meningococci

Class I outer-membrane protein and/or epitopes by

overproducing vectors either through or not through

existing open reading frames and/or manipulated

reading frames so that fusion proteins or proteins

with exchanged epitopes can be prepared.

Readily available of wild strains are:

H44/76 (B:15:P1,7.16) (Holten E., Norway, deposited

as CBS 635-89); 187 (B:4:P1,7) (Etienne J., France);

M1080 (B:1:P1,1.7) (Frasch C., USA); Swiss4

(B:4:P1,15) (Hirschel B., Switzerland); B21061

(B:4:P1,2) (Berger U., West-Germany); 395

(B:NT:P1,9) (Jónsdóttir K., Iceland); M990

(B:6:P1,6) (Frasch C., USA); 2996 (B:2b:P1,2) RIVM,

The Netherlands; M982 (B:9:P1,9) (Frasch C., USA);

53446 (B:14:P1,6) (Frasch C., USA); H355

(B:15:P1,15) (Holten E., Norway); 6557 (B:17:P1,17)

(Zollinger W., USA) and B40 (A:4:P1,10) (Achtmann M.,

West-Germany). An example of a Class 3 negative

mutant is H1115 (B:-:P1,16) deposit # CBS 636.89.

These strains were inoculated from precultures

at -70°C into shake flasks and transferred from

these into 40, 150 or 350 litre fermenter cultures.

The semisynthetic medium had the following composi-

tion: L-glutamic acid 1.3 g/l, L-cysteine.HCl 0.02

g/l, Na₂HPO₄.2H₂O 10 g/l, KCl 0.09 g/l, NaCl 6 g/l,

NH₄Cl 1.25 g/l, MgSO₄.7H₂O 0.6 g/l, glucose 5 g/l,

Fe(NO₃)₃ 100 µM, yeast dialysate.

During culturing in the fermenter, the pH and PO_2 were monitored and automatically regulated to a pH of 7.0-7.2 and an air saturation of 10%. The cells were grown to early stationary phase harvested by means of centrifuging and washing with sterile 0.1 M NaCl and stored at -20°C or freeze-dried.

EXAMPLE 2: Purification of Class 1 outer-membrane proteins from bacteriological culture

This culture can be done with the desired wild type strains, mutant meningococci strains without Class 2/3 outer-membrane proteins and/or homologous and heterologous recombinant microorganisms which express one or more of the desired meningococci Class 1 outer-membrane protein and/or epitopes by overproducing vectors either through or not through existing open reading frames and/or manipulated reading frames so that fusion proteins or proteins with exchanged epitopes can be prepared.

Readily available of wild strains are:
 H44/76 (B:15:P1,7.16) (Holten E., Norway, deposited as CBS 635-89); 187 (B:4:P1,7) (Etienne J., France); M1080 (B:1:P1,1.7) (Frasch C., USA); Swiss4 (B:4:P1,15) (Hirschel B., Switzerland); B21061 (B:4:P1,2) (Berger U., West-Germany); 395 (B:NT:P1,9) (Jonsdottir K., Iceland); M990 (B:6:P1,6) (Frasch C., USA); 2996 (B:2b:P1,2) RIVM, The Netherlands; M982 (B:9:P1,9) (Frasch C., USA); S3446 (B:14:P1,6) (Frasch C., USA); H355

(B:15:PI,15) (Holten E., Norway); 6557 (B:17:PI,17) (Zollinger W., USA) and B40 (A:4:PI,10) (Achtman M., West-Germany). An example of a Class 3 negative mutant is H115 (B:-:PI,16) deposit # CBS 636.89. These strains were inoculated from precultures at -70°C into shake flasks and transferred from these into 40, 150 or 350 litre fermenter cultures. The semisynthetic medium had the following composition: L-glutamic acid 1.3 g/l, L-cysteine.HCl 0.02 g/l, $\text{Na}_2\text{HPO}_4 \cdot 2\text{H}_2\text{O}$ 10 g/l, KCl 0.09 g/l, NaCl 6 g/l, NH_4Cl 1.25 g/l, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.6 g/l, Glucose 5 g/l, $\text{Fe}(\text{NO}_3)_3$ 100 μM , yeast dialysate. During culturing in the fermenter, the pH and PO_2 were monitored and automatically regulated to a pH of 7.0-7.2 and an air saturation of 10%. The cells were grown to early stationary phase harvested by means of centrifuging and washing with sterile 0.1 M NaCl and stored at -20°C or freeze-dried. The bacterial mass was for example extracted with the aid of 0.5 M CaCl_2 , 1% (w/v) Zwittergent 3-14 (Zw 3-14) and 0.14 M NaCl, pH 4.0, using 100 ml per gram of freeze-dried bacterial mass. The suspension was stirred for 1 hour at room temperature and then centrifuged (1 hour, 3000 x g), after which the supernatant was collected in a sterile manner. 20% ethanol (v/v) was added to the supernatant and after stirring for 30 min. the product was centrifuged (30 min., 10,000 x g), after which the supernatant was collected aseptically. The supernatant was then concentrated by means of

Cyanogen bromide was used to prepare fragments of meningococci Class 1 outer-membrane proteins. The purified Class 1 or mixtures of Class 1 or 3 outer-membrane proteins were taken up in 70% (v/v) formic acid and treated with a 10-fold excess of CNBr for 16 hours at room temperature. The CNBr and the formic acid were removed by means of evaporation

EXAMPLE 3: Preparation and Characterization of Class 1 OMP Peptide Fragments

dialfiltration in an Amicon Hollow Fiber System (HID x 50, cut off 50,000) and CaCl_2 and ethanol were removed. The concentrate was diluted with 0.1 M sodium acetate, 25 mM EDTA, 0.05% ZW 3-14 having a pH of 6.0 to the original volume and then concentrated again by means of dialfiltration. This procedure was repeated five times. The pH of the final concentrate was adjusted to a value of 4.0. 20% (v/v) ethanol was added to the concentrate and, after stirring for 30 min., the product was centrifuged (30 min., 10,000 x g). The whole proteins are purified with the aid of column chromatography in the presence of detergent, for example ZW 3-14. Often gel filtration over Sephacryl S-300 as well as the ion exchange over DEAE Sepharose is applied (Beuvery et al. (1986) supra). The used extraction method, detergents, column chromatography are not the only applicable method yet only serve as examples and must not be regarded as restrictive.

-38-

and replaced by 0.2 M Tris.HCl, 6 M urea solution, pH 7.2. The supernatant was prepurified by means of gel filtration over Sepharil S-200 and subsequently purified with the aid of TSK-2000 gel filtration via HPLC. Beuvery et al., (1986) *supra*.

Enzymatic digestion of CB2 fragments

To further delineate the epitopes, the meningococcal CB2 fragment was subjected to digestion with EndoARG-C, EndoGLN-C or V-8 and the resulting fragments isolated by HPLC. Briefly, 20 nmoles of CB2 fragment in 1 ml of 25 mM phosphate/0.1 mM tris buffer (pH 8.0) containing 3M urea was digested at 37°C with 0.2 nmoles of EndoARG-C (1mg/ml in distilled water) or 0.22 nmoles of EndoGLN-C or V-8 (1 mg/ml in distilled water) for 14-18 hours. The resulting digested fragments were separated by reverse phase HPLC using a Vydac-C4 column and a trifluoroacetic acid-acetonitrile gradient. The main peak eluted from the EndoARG-C digestion had an apparent molecular weight of 7-9 Kdal while the main peak observed following EndoGLN-C or V-8 had an apparent molecular weight of 4-6 Kdals. The isolated peaks were subsequently shown by Western blot to react to a pool of monoclonal antibodies (Adam I, 62-D12-8, MN5-C11G and MN14-C116).

The P1.16 epitope appears to be present on the

C-terminal CNBr fragment of the Class I outer-mem-

brane protein of strain H44/76 (B:15; P1.7.16).

Further characterization of the P1.16 epitope was

carried out through amino acid sequence determina-

tion of the 17Kd (N-terminal) and 25Kd (C-terminal)

CNBr fragments. The C-terminal 25Kd is further

fragmented with V8 protease, endoLysC, endoGlu-C and

endoArg-C. Fragments which were positive with the

P1.16 monoclonal antibody were sequenced as far as

possible. The sequences which were obtained are as

follows:

N-terminus of whole protein:

DVSLYGEIKAGVEDRNNYQLQLTEAQUAAGN...

N-terminus of 25Kd C-terminal CNBr

fragment : (M) PVSVRVDSPEFSGSVQFVPIONS-

KSAYTPAYYTKDTNNN...

Fragments which react with P1.16 monoclonal anti-

bodies were isolated using V8 protease and endoArg-C

fragmentation with a molecular weight of 7-9Kd and

4-6Kd respectively. The N-terminal sequences hereof

are as follows:

V8 7-9Kd fragment: FSGFSGSVQFVPIQNSKSAYTPAYYTKDTN...

Arg-C 4-6kd fragment: PVSVRKYDSEPFSGFSGVQFVPI-
QNSKSAVTPAVYTK...

EXAMPLE 4: DNA Sequences of Class I OMP Genes

Amino acid sequences of Class I OMP were deduced from the nucleotide sequence of the structural genes of four meningococci Class I OMP's with various subtypes. Comparison with four amino acid sequences enabled a prediction of the composition and the location of these epitopes. Further, the aid of peptide synthesis and the demonstration of binding of the respective monoclonal antibodies. Class I OMP genes were cloned into lambda gt11 (as described for P1, 16 in Barlow et al., (1987) Infect. Immun. 55: 2743-2740) and subcloned in M13 sequencing vectors and the DNA sequence was determined by standard chain termination dideoxynucleotide techniques. The complete derived amino acid sequence for P1, 16; P1, 15; P1, 7.16; and P1, 2 proteins are as follows:

P1.16:	DVSLYGEIKAGVEGRNIQALTEPQVTNGVGNQV--KVTAKSRIRTKIS *****	10
P1.15:	DVSLYGEIKAGVEGRNFQLTLEPP-SKSQF--QV--KVTAKSRIRTKIS *****	20
P1.7.16:	DVSLYGEIKAGVEGRNYQLTEAQAANGASGQVKVTKVTKAKSRIRTKIS *****	30
P1.2:	DVSLYGEIKAGVEGRNIQALTEPQVTNGVGNQV--KVTAKSRIRTKIS *****	40
		50

 S+
 AYTPAHFVQOTFQPTLVPAVVGKPGSDVYYAGLNYKNGGFAAGNYAFKYAKHNAVGRDAF

 AYTPAYYTKNTNNLTLVPAVVGKPGSDVYYAGLNYKNGGFAAGNYAFKYARHNAVGRDAF

 AYTPAHYTRQNNNTDV-FVPAVVGKPGSDVYYAGLNYKNGGFAAGNYAFKYARHNAVGRDAF

 AYKPAYYTKNTNNLTLVPAVVGKPGSDVYYAGLNYKNGGFAAGNYAFKYARHNAVGRDAF

 180 190 200 210 220 230

 NOFDASKAIDPMDSNVVAQGLGIFKRMDMPVSVRYDSPFSGSVQFVPAQNSKS

 NOFDASQAIDPMDSNNDVAQGLGIFKRHDMVSVRYDSPFSGSVQFVPIQNSKS

 NOFDASQAIDPMDSNNDVAQGLGIFKRHDMVSVRYDSPFSGSVQFVPIQNSKS

 NOFDASQAIDPMDSNNDVAQGLGIFKRHDMVSVRYDSPFSGSVQFVPAQNSKS

 120 130 140 150 160 170

 DGSFIGFKGSEDLGEGTLKAVWQLEQDVSAAGGATQWGNRESFVGLAGEFGTLRAGRA

 DGSFIGFKGSEDLGEGTLKAVWQLEQDVSAAGGATQWGNRESFVGLAGEFGTLRAGRA

 DGSFIGFKGSEDLGEGTLKAVWQLEQDVSAAGGATQWGNRESFVGLAGEFGTLRAGRA

 DGSFIGFKGSEDLGEGTLKAVWQLEQDVSAAGGATQWGNRESFVGLAGEFGTLRAGRA

 60 70 80 90 100 110

290 280 270 260 250 240
ELFLIGS-DEAKGTDPLKNNHQBRLTGGEYEEGGLNLAQAQDLSENAD--KTKNSTT
ELFLIGS-DEAKGTDPLKNNHQBRLTGGEYEEGGLNLAQAQDLSENAD--KTKNSTT
ELFLIGS-DEAKGTDPLKNNHQBRLTGGEYEEGGLNLAQAQDLSENAD--KTKNSTT
ELFLIGS-DEAKGTDPLKNNHQBRLTGGEYEEGGLNLAQAQDLSENAD--KTKNSTT

350 340 330 320 310 300
EIAATASRYRFGNAVPRISSYAHGFDFIERGKKGENTSYDQIIAGVDYDFSKRTSAIVSGAM
EIAATASRYRFGNAVPRISSYAHGFDFIERGKKGENTSYDQIIAGVDYDFSKRTSAIVSGAM
EIAATASRYRFGNAVPRISSYAHGFDFIERGKKGENTSYDQIIAGVDYDFSKRTSAIVSGAM
EIAATASRYRFGNAVPRISSYAHGFDFIERGKKGENTSYDQIIAGVDYDFSKRTSAIVSGAM

370 360
LKRNTGIGNYTIQINAASVGLRHKF
LKRNTGIGNYTIQINAASVGLRHKF
LKRNTGIGNYTIQINAASVGLRHKF
LKRNTGIGNYTIQINAASVGLRHKF

+ Note this amino acid 15 is located between A.A.S.184 and 185 of this sequence

EXAMPLE 5: DNA Sequencing of Class I OMP Genes from different *N. meningitidis* Serotypes

The Polymerase Chain Reaction (PCR) technique of Mullis and Faloona (Methods in Enzymol. 155:335-50, 1987) was used to amplify the entire Class I OMP gene and specific fragments according to the scheme shown in Figure 1.

Primers were synthesized on an Applied Biosystems 380B DNA synthesizer and used in standard PCR 30 cycle amplification reactions using Taq polymerase in a Thermal Cycler (Perkin-Elmer Cetus, Norwalk, CT) according to the recommendations of the Supplier. Amplified fragments of about 1300, 900 and 450bp were generated from each serosubtype genomic DNA preparation from the primer combinations shown in Figure 1. The primers used had the following sequences:

- PR1: (41 bases with universal primer extension)
TGT AAA ACG ACG GCC AGT TTG AAG ACG TAT CGG GCG TTT GC
- PR2: (42 bases with universal primer extension)
TGT AAA ACG ACG GCC AGT GGC GAA TTC GGT ACG CTG GCG GCC
- PR3: (42 bases with universal primer extension)
TGT AAA ACG ACG GCC AGT CAT CAG GTA CAC GCG CTG ACG GCG
- PR4: (40 bases with universal primer extension)

-44-

TGT AAA ACG ACG GCC AGT GCA GAT TGG CAG TCA GAT TGC A

PR5: (40 bases with universal primer extension)
TGT AAA ACG ACG GCC AGT GCG ATC GGT ACC TTT GGC TTG A

PR6: (40 bases with universal primer extension)
TGT AAA ACG ACG GCC AGT AAC TGA TTC GCA ACG CGA CCG G

FWD: (24 bases)
TTG AAG GAC GTA TCG GGT GTT TCG

REV: (23 bases)
GCA GAT TGG CAG TCA GAT TGC TT

Excess single stranded template for sequencing

was synthesized in an 'asymmetric PCR' amplification using 100x excess of primer carrying an 18 base

extension at the 5' end corresponding to the universal fluorescent sequencing primers used with an

Model 370A Automated DNA Sequencer (Applied Biosystems, Foster City, CA). Taq polymerase was used in

a standard dideoxynucleotide chain termination sequencing reaction with the PCR generated single

stranded Class I gene fragments as templates. Derived sequences for gene segments of strains

H44/76 (PI.7.16), M1080 (PI.1.7), H355 (PI.15), 6940 (PI.6), 6557 (PI.14), 870227 (PI.10) and B40 (PI.10)

are shown in Figures 2 and 3.

EXAMPLE 6: Confirmation of Amino Acid Sequences of
Class I OMP Subtype Epitopes

From these gene sequences confirmed by direct sequencing of Class I OMP genes, it was deduced that the sequences corresponding to amino acids 24-34 and 176-187 of P1.16 are markedly variable in the four Class I OMP sequences. Three amino acid sequences N-terminal or C-terminal from these positions should also be considered for possible inclusion in these epitopes to allow for maximizing epitope stability presentation and unexpected insertions or deletions in the native protein sequence. Further the DNA and amino acid sequences of other Class I OMPs should be compared with the P1.7,16 sequence to allow for maximum alignment and epitope prediction. The first variable region epitope and second variable region epitope are called VR1 and VR2 respectively. These regions encode the subtype epitopes as was confirmed with the aid of peptide synthesis and the reaction of the peptides with P1.2; P1.7; P1.15 and P1.16 specific monoclonal antibodies. A complete set of overlapping decapptides staggered by 5 amino acids were prepared using the P1.16 protein sequence. The anti-P1.16 monoclonal antibody reacted with the decapptide YTKDITNNNL from P1.16 reacted as expected and no other decapptide. (Figure 4). Of overlapping decapptides provided with a one (1) amino acid sequence shift in the region 24-34

and 176-187 of the Class I OMP of strains H44/76

(P1.7,16), MC50 (P1.16) and MC51 (P1.15) more than one peptide reacted with the subtype specific

monoclonal antibody. In most cases one or more of the group of these overlapping peptides reacted with

the subtype specific monoclonal antibody more strongly than others (Figure 5).

These peptides are designated as the VR1 and

VR2 epitopes. In the P1.7,16 strain, the sequence

YYTKNTNNL is present, the change D to N at residue

180 does have some effect on reducing antibody bind-

ing. The sequence HYTRQNNNTDVF in P1.15 in the same

relative position in the protein as the P1.16

epitope and is responsible for binding to the

anti-P1.15 monoclonal antibody. AQVANGCASC shows

some binding and peptides 1-3 amino acids downstream

show far greater binding to the P1.7 monoclonal

antibody. Sequence HFVQQTTPQSQP of VR2 is

responsible for binding to the anti-P1.2 monoclonal

antibody. It is probable that the sequences

QPQVTNGVQGN and PPSKSP in the P1.16 and P1.15

proteins also represent epitopes.

Example 6B: Class I OMP Constant Region Epitope

Identification

Peptides forming surface loops were prepared and conjugated to tetanus toxoid. A Biolynx 4170 automated peptide synthesizer (Pharmacia/LKB) was used for continuous flow solid-phase synthesis with

the following exception. In the last cycle of the synthesis SAMMA-OFIP (0.5 mmol) (Drijfhout, J.W. (1989), Ph.D. Thesis, Leiden, The Netherlands) was coupled in the presence of 1-hydroxybenzotriazole (0.5 mmol) for 30 min., using a standard protocol with omission of the piperidine-treatment (i.e. the "Fmoc-deblocking step" which in this case would cause undesirable S-deacetylation). These are referred to as SAMMA-peptides.

The peptides and their surface region location which were conjugated to IT are as follows:

Name	Peptide	Region
LBV 017	176 185 XGGYTKDITNNIL	P1.16, loop 4
018	24 33 XGGAQAAAGCAGS	P1.7, loop 1
024	276 291 XGGLSENGDKAKTKNSTTE	P1.16, loop 6
025a	245 XGGAFAELFLIGSATSDAAG	P1.16, loop 5
025b	223 XANVGRAFAELFLIGSATSDAAG	P1.16, loop 5
026	124 137 XGDSNNDDVASQLQIFK	P1.16, loop 3
027	XADLNTDAERVAVNATMAHFV	Class 2, loop 5
028a	329 XGGGKKGENTSYDQ	Class 1, loop 7
028b	317 XGGERGKRGENTSYDQ	Class 1, loop 7
029	XGGVKDAGTYRAQGGKSKTATQ	Class 2, loop 1
030	78 90 XGGSVAEGCAGQVGN	P1.16, loop 2
031	352 366 XKRNTGIGNYTIQINAA	P1.16, loop 8
032	16 34 XGGNIQAQLTEQPVITNGVGN	P1.16, loop 1

Conjugation of SAMa-peptides to tetanus toxoid was performed as follows. A solution of N-succinimidyl bromoacetate (4.7 mg, 10 μ mol) in DMF (100 μ l) was mixed with a solution of tetanus toxoid (100 μ l) in 0.1 M sodium phosphate buffer pH. (TT) (20 mg) in 0.1 M. After 1h, 1.8 ml of the reaction mixture was subjected to gel filtration using a Sephadex PD-10 column (Pharmacia) equilibrated in 0.1 M sodium phosphate, containing 5mM EDTA (pH buffer) pH 6.1. The bromoacetylated tetanus toxoid was eluted with the same buffer and collected in 3.5 ml. The solution of bromoacetylated tetanus toxoid (1.2 ml) was added to the SAMa peptide (4.5 mg, 3 μ mol) and degassed with helium. Next, 150 μ l of 0.2 M hydroxylamine (in PE buffer, pH 6.1) was added. After 16 h remaining bromoacetyl groups were blocked by addition of 2-aminoethanethiol hydrochloride (4 μ mol) in buffer, pH 6.1 (150 μ l). After a further period of 16 h, the peptide-TT conjugate was purified by gel filtration over a PD-10 column using PE buffer, pH 6.1, as the eluant. The appropriate fractions were combined and stored at 4°C.

To determine the immunological activity, 25 μ g (total protein) per dose of a peptide-TT conjugate was injected subcutaneously at weeks 0 and 4 into 6-8 week old NIH outbred mice. (Note: Vaccine LBV 017-TT and LBV 018-TT were used at 10 μ g total protein/dose.) Sera were collected 6 weeks following the first dose and evaluated for antibody

The results are presented in Table 2 below.

(Poolman, J.T. et al., (1985) supra.)
Bactericidal activity (BC) of sera was also measured.
Immun. 57:1005 and the unconjugated peptides.
I OMP (Poolman, J.T. et al., (1989) Infect. and
wells: Outer membrane protein (OMP), purified Class
following antigens were coated into the microtiter
(1983) Infect. and Immun. 40:3690380). The
response in an ELISA assay (Beuvery, E.C. et al.

TABLE 2

Bactericidal	Test	Synth. Peptide	Class 1 OMP	OMC	Vaccine	

<1:64	ND	ND	1:2700	1:900 (0.05)*	LBV 018-TT	LBV 018-TT
<1:64	ND	ND	1:900	1:900 (1)	LBV 017-TT	LBV 017-TT
<1:64		1:900 (homol.)	1:100	1:100	LBV 024-TT	LBV 024-TT
<1:64		1:2700 (homol.)	1:100	-	LBV 025a-TT	LBV 025a-TT
<1:64		1:8100 (homol.)	1:300	1:2700 (4)	LBV 025b-TT	LBV 025b-TT
<1:64		- (homol.)	-	-	LBV 026-TT	LBV 026-TT
<1:64		1:300 (homol.)	1:300	-	LBV 027-TT	LBV 027-TT
<1:64		1:2700 (homol.)	-	1:100	LBV 028a-TT	LBV 028a-TT
<1:64		1:900 (homol.)	1:100	1:100	LBV 028b-TT	LBV 028b-TT
<1:64		1:8100 (homol.)	1:100	-	LBV 029-TT	LBV 029-TT
<1:64		1:2700 (homol.)	1:100	-	LBV 030-TT	LBV 030-TT
<1:64		- (homol.)	1:100	-	LBV 031-TT	LBV 031-TT
<1:64		1:900 (homol.)	1:100	-	LBV 032-TT	LBV 032-TT

* numbers in () indicate O.D. level showing this titer

These data suggest that of the constant surface loops tested of Class 1 and 2 OMPs of *N. meningitidis* loop 5 appears to represent at least one region that will produce antibodies which will cross-react with Class 1 and Class 2 OMP of many strains of *N. meningitidis*.

EXAMPLE 7: Construction of recombinant flagellins
expressing meningococcal epitopes

To create hybrid flagella containing epitopes from class I meningococcal epitopes, a series of oligonucleotides was designed based on primary protein sequence data and epitope mapping data. Two oligonucleotides based on VR1 or VR2 epitopes of outer membrane P1.7.16 were designed so that they could be cloned in single or multiple copies into a cloning region within the gene for *S. muenchen* flagellin. Translation termination signals were included on the non-coding strand of the oligonucleotide to facilitate screening by expression of the cloned inserts.

The plasmid vector pPX1650 containing the entire coding region and promoter regions for the structural gene for flagellin H1-d of *Salmonella muenchen* (deposited at the ATCC, accession #67685) was modified to contain several unique cloning sites suitable for the insertion of either oligonucleotides or gene fragments in each of the three reading frames of the flagellin gene (Figure 6). First, pPX1650 was digested with *EcoRV*, which cleaves pPX1650 twice, 48 base pairs apart, and religated to yield a plasmid, pPX1651, which has a unique *EcoRV* cloning site and which results in a 16 amino acid deletion in the flagellin protein. pPX1651 was identified by screening *E. coli* recombinants on Western blots probed with polyclonal antibody

directed against HI-d flagellin. pPX1651 was identified amongst several candidates having flagellins smaller than wild type flagellin (of 1650) and was verified by sequencing. Second, pPX1651 was restricted with BamHI and religated after filling out the overhanging ends with Klenow enzyme to remove the unique BamHI restriction enzyme site in the polylinker region of the vector. As a final step, the resulting vector was digested with EcoRV and the following oligonucleotide linker was inserted:

```

5'      ATG ATC GAT GCA TTC      3'
3'      TAG TAG CTA CTA AAG      5'

```

Candidates were screened for the newly created BamHI sites and several candidates having BamHI sites were screened for orientation of the linker by double strand DNA sequencing methodology. One candidate having the linker in the above orientation was retained as pPX1647:

```

5' . . . . GAT ATC ATC GAT GCA TTC ATC . . . .
          EcoRV      ClaI      BamHI

```

Plasmid pPX1647 (Figure 7) was digested with BamHI and either oligonucleotides for VR1 or VR2 were cloned into E. coli cells. Screening for desired recombinants was accomplished by digesting plasmid miniplysate DNA with appropriate diagnostic

restriction enzymes and screening for expression by probing hybrid flagella for decreased mobility on SDS-PAGE gels with specific flagellar antiserum (H1-d). A number of the resultant clones showed decreased mobility on SDS-PAGE, indicating proper insertion of one or more of the oligonucleotides for VR1 or VR2. Several of each were retained for analysis by DNA sequencing. Clone CBI-2 results from tandem insertion of two copies of the VR1 oligonucleotide and clone CBI-4 results from insertion of four oligonucleotides. Likewise CB2 P contained a single insert of the VR2 oligonucleotide and CB2 W showed the expected trimetric insert, CB2 P clone contained a single base pair change which resulted in a change from Leu to Phe in the expressed VR2 fusion protein and was not retained for further study. The recombinant flagellin clones in E. coli were probed with monoclonal antibodies (Abdillahi and Poolman, Microbiol. Pathogenesis 4:27-32, 1988; RIVM, The Netherlands) known to react with either VR1 or VR2 epitopes. Monoclonals Adam-1 (P1.7) and Mnl4-C11-6 (P1.7) react with hybrid flagellin containing 2 or 4 tandem inserts of VR1, but do not react with clones containing VR2. The weaker reaction of both monoclonals with CBI-2 than with CBI-4 is likely due to epitope density. By the same token, monoclonals 62 (P1.16) and Mnl5-c11-G (P1.16) react with CB2 W clone, but not with the VR1 inserts. The CB2 P clone fails to react with either VR2 antibody, probably due to the Leu to Phe change.

Each of these clones was transformed into an *Escherichia coli* strain (SL5927), having a Tn10 insertion in the H1-d locus, to examine the functioning of the hybrid flagella. Each of the four clones resulted in motile bacteria; motility of the transformants was inhibited by the corresponding monoclonal antibody, including clone CB2 F, indicating affinity of the VR2 monoclonal for the epitope in intact flagella. This result indicates that epitopes are exposed at the cell surface and are accessible to antibody.

Hybrid flagellin containing both VR1 and VR2 epitopes were created by cleaving either CB1-2, CB1-4, or CB2 with $\overline{\text{BamHI}}$ and cloning the heterologous epitope. Clones CB12-7 and CB12-10 result from the in-frame insertion of a single copy of the VR2 oligonucleotide behind either 2 or 4 VR1 tandem inserts, respectively; clone CB21-F arose from the insertion of one copy of the VR1 epitope behind 3 tandem copies of VR2. CB12-7 and CB12-10 are recognized only by VR1 monoclonal antibody and CB21-F is recognized only by VR2 monoclonal. These results, taken together with DNA analysis revealing predicted sequences, indicate epitope density is too low in the combined hybrids. To create a hybrid flagellin with increased density of both VR1 and VR2 epitopes, CB12-10 was digested with $\overline{\text{BamHI}}$ and VR2 encoding oligonucleotides were inserted. Clone 12-10-6 contains two further tandem inserts of the VR2 epitope, resulting in a hybrid flagellin

molecule in which four tandem copies of VR1 are followed by three copies of VR2. As is shown in Figure 3a and b, three of the hybrid flagellin vaccine candidates have the expected molecular properties. The flagellin (pCB1 x 4) containing 4 copies of VR1 reacts with anti-H1-d (anti-flagellin) and anti-VR1 monoclonal antibodies, but not with anti-VR2 monoclonal antibodies; the flagellin (pCB2-W) containing 3 tandem copies of VR2 reacts with anti-H1-d and anti-VR2 antibodies, but not with anti-VR1; the combined hybrid containing copies of VR1 and 3 copies of VR2 reacts with both anti-VR1 and anti-VR2 monoclonal antibodies. The combined hybrid specified motility when introduced into a non-motile recipient *S. dublin* strain. As a subunit vaccine, the goal is to obtain suitable initial vaccine candidates in high quantity and high purity. A suitable vaccine candidate can be chosen from the above type constructions based on reactivity to monoclonal antibodies and function of flagella in non-motile *Salmonella* host strains. A subunit flagellin vaccine may not need to retain all functional aspects of a parental flagellin, but should at least retain surface localization for purification purposes. Several subunit flagellin meningococcal vaccine were chosen from the above described hybrid molecules based on reactivity to monoclonal antibodies and implied surface localization based on restoration of bacterial motility.

The three hybrid flagellin vaccine candidates and a wild type (derived from PFX1650) were inoculated into four-liter baffled Fernbach flasks containing 1 liter of LB broth. Bacterial cultures were incubated at 37°C with shaking (200 rpm) for 22-24 hr. Under these conditions of culturing, the bulk of the flagella were sloughed from the bacterial cell surface and were localized in the supernatant culture medium. To obtain suitable material, flagella were isolated from 6-8 liters of culture medium. To obtain purified flagellin preparations for vaccination studies, flagellar filaments were harvested from bacterial culture supernatants by the following procedure: Ammonium sulfate was added to culture supernatant so that final solution was 50% saturated; the solution was stirred gently at 4°C for several hours and the precipitated material was collected by centrifugation in a GSA rotor at 5000

EXAMPLE 8: Initial Purification of recombinant flagellin molecules

Three flagellin vaccine candidates contained either 4 tandem inserts of VR1, 3 tandem inserts of VR2, or 4 VR1 inserts followed by 3 VR2 inserts. Because flagellin is a major protein of *Salmonella*, it is possible to easily purify sufficient material for vaccination studies using techniques established for flagella purification (Logan et al., J. Bacteriol. 169: 5072-5077, 1987).

To prepare highly purified flagellins, *Salmonella* expressing the constructions, in particular PCB12-10-6, was grown as described above and the cells pelleted at 10,000G. The culture supernatant was then precipitated with 50% ammonium sulfate, centrifuged at 10,000G and resuspend in 30 ml PBS. The resuspended pellet was dialyzed against 10mM Tris buffer (pH-8.0) containing 6M urea, 1mM PMSF, 2mM NEM, and 5mM EDTA overnight at 4°C. Dialyzed material was then passed over two DEAE sepharose minicolumns (3.0 ml volume, 4.0 ml eluent over each). The columns were eluted (5X) with 50mM NaCl in 10mM Tris (pH-8.0) containing 6M urea and then with 1M NaCl in 10mM Tris (pH-8.0) containing 6M urea. The first four elution collections (20 ml) of urea. The 50mM NaCl were pooled and dialyzed against 1.0 liter 10mM acetate buffer (pH-4.0) in 6M urea at

EXAMPLE 9: HPLC Purification of recombinant flagellins

The following method. flagellin, was subjected to further purification by pelleted material, which consisted primarily of rotor to pellet the flagellar filaments. The dialyzed material was subjected to high speed centrifugation at 100,000 x g for 1 hour in an SW-27 dialyzed against PBS at 4°C for 12-15 hrs. The precipitated material was reconstituted in PBS and rpm for 30 minutes. The collected ammonium-sulfate

Group C meningococcal capsular polysaccharide (GCM CPS: lot # 86 NM 01) was prepared essentially according to Bunde et al., J. Biol. Chem. 249: 4797-801, 1974). *Neisseria meningitidis* strain C11 was obtained from the Walter Reed Army Institute (Washington, DC). The strain was precultured twice on sheep blood agar plates, then used for the inoculation of a liquid seed culture medium *Neisseria* chemically defined medium, (NCDM) Kenney et al., Bull. W.H.O.

EXAMPLE 10: Preparation of meningococcal-flagellin glycoconjugate

The dialyzed fractions were then loaded onto a TSK SP PW cation exchange HPLC column (75mm x 300mm). The column was eluted with a mobile phase consisting of 10mM Acetate (pH=4.0) containing 6M urea. A gradient 0 - 300mM NaCl was established in 10mM acetate (pH=4.0) containing 6M urea over the 5 - 30 min interval. After 30 mins the gradient went from 300mM to 1M NaCl in 10mM acetate in 6M urea over the next 5 min. The flagellin construct was collected at approximately 24 min. which corresponds to about 200mM NaCl. The fraction was dialyzed against PBS and purity determined on the material was established by Western blots using anti-flagellin antibody. A representative HPLC analysis and SDS-PAGE are shown in figures 8 and 9 respectively.

37 469-73, 1967). Finally, 40 l of liquid medium (NCDM) in a fermentor was inoculated with the liquid preculture. The purity of the strain was checked at each stage. After centrifugation, the supernatant was precipitated by addition of cetylal to a final concentration of 0.1%, and the insoluble complex re-dissolved in cold 1 M calcium chloride (CaCl_2) (Gotschlich et al., J. Exp. Med. 129:1349-65, 1969). Ethanol (96%) was added to a final concentration of 25% (v/v). After 1 h, the suspension was centrifuged (1 h, 50,000 g), the supernatant was collected, and its ethanol concentration was increased to 80% (v/v). After 1 h, centrifugation (20 min, 5,000 g) yielded a precipitate which was washed successively with absolute ethanol, acetone, and diethyl ether, and then dried in a vacuum desiccator over phosphorus pentoxide (P_2O_5) to constant weight. This crude CPS was stored at -20°C.

In order to obtain a purer preparation, the CPS

was then dissolved in sodium acetate buffer (1.10 dilution of a saturated solution, pH 7.0) and extracted four times with hot phenol (Westphal et al., Z. Naturforsch. 7b:148-55, 1952). After dialysis of the combined aqueous phases against 0.1 M CaCl_2 , followed by centrifugation (3-5 h, 100,000 g), a final ethanol precipitation was performed on the clear supernatant, and the resulting precipitate washed with organic solvents

and dried, as described above. The pure CPS was then stored at -20°C.

At each stage of the purification process, the CPS was analyzed for carbohydrate N-acetyl-neuraminic acid, NANA) (Svennerhold, *Biochim. Biophys. Acta* 24:604, 957), O-acetyl (Hestrin, *J. Biol. Chem.* 180:249, 1949), and protein (260 nm detection) content, and its molecular weight checked by gel filtration.

Group C meningococcal capsular polysaccharide (GCM CPS) was simultaneously depolymerized and activated via sodium periodate (NaIO_4) oxidation in aqueous buffer (Anderson et al., *J. Immunol.* 137:1181-6, 1986; Eby et al., *Pediatr. Res.* 20:308A, 1986, Anderson et al., *J. Pediatr.* 111(5):644-50, 1987; Anderson, U.S. Pat. 4,762,713; 1988). The reaction was monitored by high performance gel permeation chromatography (HPLC) in aqueous eluent, using ultraviolet (UV) and refractive index (RI) detection. The reaction was stopped and the activated oligosaccharides (GCM OS) were desalted by low pressure gel permeation (GPC) in water, and then lyophilized. A solution was then prepared in water and subsequently frozen for temporary storage. GCM OS and flagellin PCB12-10-6 were mixed in aqueous neutral buffer and the conjugation was initiated by addition of sodium cyanoborohydride (NaBH_3CN) (Anderson, U.S. Patent 4,762,713, 1988; U.S. Patent 4,673,574, 1987; U.S. Patent 4,761,283, 1988). The reaction was carried out for 5 days, while being

monitored by HPGPC. It was finally stopped by dialysis/concentration on centrifugal microconcentrators. The final preparation was stored in the cold, in the presence of thimerosal to prevent bacterial growth. The resulting glycoconjugate not only provides a mechanism to present the expressed VRI and VR2 meningococcal epitopes to the immune system but also serves as a carrier molecule for the presentation of a meningococcal oligosaccharide. In preparation of the conjugate, the following conditions were employed. Purified flagellin PCB12-10-6 was dissolved in 15% sucrose (3.5 mg/ml) and then stored at -20°C. GCM CPS (9.7 mg; final concentration: 5 mg/ml) was oxidized by 100 mM NaIO₄ in 0.05 M sodium phosphate buffer (pH 6.2 - 6.5) at RT, in the dark, with agitation. Aliquots (100 µl) were withdrawn at regular intervals, the reaction stopped by addition of ethylene glycol (10 µl), and analysis was performed by HPGPC on Waters (Milford, MA) UltrahydrogelTM 250 + 120 (2 columns coupled; 2 x 300 mm x 7.8 mm) in 0.2 M phosphate-saline buffer (PBS; 0.2 M sodium phosphate, 0.9% NaCl, pH 7.8), at a flow rate of 0.8 ml/min, using UV (206 nm) and RI detection. After 2 h 30 min, the reaction was stopped by addition of ethylene glycol (1/10 of the reaction volume), and the GCM OS were desalted by GPC on Bio-Rad (Richmond, CA) Bio-Gel^R P-2 (200-400 mesh, 30 cm x 1.5 cm) in water, at about 18 ml/h. Fractions were collected (1.2 ml) and analyzed for the presence of NANA the carbohydrate N-acetyl-

-63-

neuraminic acid (NANA) (Barry *et al.*, J. Gen. Microbiol. 29:335-52, 1962) and aldehydes (Porro *et al.*, Anal. Biochem 118:301-306, 1981). Positive fractions were pooled and lyophilized. Desalted GCM OS (4.7 mg) were then dissolved in water (10 mg/ml) and frozen at -20°C.

Both GCM OS and PCB12-10-6 solutions were analyzed by HPGPC (UV at 206 and 280 nm respectively) before being frozen, and prior to the conjugation. No degradation occurred during storage, as ascertained by the exact similarity of the elution profiles.

GCM OS (2 mg; final concentration: 2.6 mg/ml) and flagellin PCB12-10-6 (2.3 mg; final concentration: 3 mg/ml) were mixed in a polypropylene tube in 0.4 M sodium phosphate buffer (pH 7.0), and NaBH_3CN was added (12 μmoles) to initiate the conjugation (Anderson, U.S. Patent 4,762,713, 1988; U.S. Patent 4,673,574; U.S. Patent 4,761,283). The reaction mixture was left one day at RT, then 4 days at 35°C, without agitation. The reaction was monitored by HPGPC (UV at 280 nm) at different stages, and finally stopped by dialysis/concentration on microconcentrators. The final preparation was analyzed for NANA (Barry *et al.*, J. Gen. Microbiol. 29:335-353, 1962) (0.09 mg at 0.12 mg/ml) and protein (Lowry *et al.*, J. Biol. Chem. 193:265-275, 1951) (1.12 mg; 1.45 mg/ml) content. It was then stored at 4°C in the presence of thimerosal (0.01%, w/v) to prevent bacterial growth.

Peptides designated as M20 and M21 were produced on an ABI model-peptide synthesizer by solid phase synthesis using the tBoc chemistry were coupled to CRM¹⁹⁷ (prepared as described by Andersen, U.S. Patent No. 4,762,713) using a bifunctional crosslinking agent, sulfosuccinimidyl (4-iodoacetate) amino benzoate (Sulfo SIAB; purchased from Pierce) following the modification of a published procedure (Weltman, J.K. et al., (1983) Bio Techniques 1, 148-152). Briefly CRM¹⁹⁷ was activated by sulfo SIAB resulting in the formation of an amide bond between SIAB and amino groups of CRM¹⁹⁷. After the removal of unreacted crosslinker from the activated CRM¹⁹⁷ by gel filtration, peptide (M20 or M21) containing linking spacer (represented in underlined letters) with carboxy terminal cysteine residue was

EXAMPLE 11: Conjugation of Meningococcal peptides to CRM and bovine serum albumin

The conjugate preparation was also checked by SDS-PAGE (silver nitrate stain) and Western blots analyzes. Several high molecular weight bands appeared on the gel above the pure PCB12-10-6 band and near the stacking well, the latter being an evidence that cross-linking occurred during conjugation. Western blot analyzes showed that each band was reactive with the antisera used (anti-GCM, -VR1, and -VR2), proving covalency of the conjugate bonds.

-65-

mixed with activated CRM and incubated at room temperature for 2-4 hours. Following the reaction, the conjugated material was dialyzed extensively against PBS at 4 C.

The sequence of M20 peptide (VR2 epitope) is as follows:

H-Tyr-Tyr-Thr-Lys-Asp-Thr-Asn-Asn-Asn-Leu-Thr-Leu-V-
al-Pro-Ala-Gly-Ala-Cys-OH

The sequence of M21 (VR1 epitope) peptide is:

H-Ala-Gln-Ala-Ala-Asn-Gly-Gly-Ala-Ser-Gly-Gln-Val-L-
ys-Ala-Gly-Ala-Cys-OH.

Conjugated materials were subjected to SDS PAGE, transferred to PVDF membranes (Immobilon, Millipore) and reacted with specific monoclonals which recognize VR1 and VR2 epitopes. Figure 10a and 10b show the western blot analysis of M20 and M21 CRM¹⁹⁷ conjugates, against a pool of VR1 and VR2 specific monoclonals (Adam I, G2-D12-8 (P1.7), MNS-G11-G (P1.16) and MN14-G11-6 (P1.7)).

In order to assay the antibody response to M20 and M21 peptide by enzyme linked immunosassay procedure, BSA conjugates were prepared by using a different bifunctional crosslinking agent, N-Succinimidyl Bromoacetate as described by Bernatowicz and Matsueda (Anal. Biochem. 155, 95-102 (1986)).

Covalent coupling of peptide to the protein was confirmed by western blotting of electrophoresed samples as described for CRM¹⁹⁷ conjugates.

EXAMPLE 12: Retention of T cell activity by M20

and M21-CRM¹⁹⁷ conjugates

To determine whether conjugation of the VRI and VR2 epitopes to CRM¹⁹⁷ adversely affect the T cell recognition of the CRM¹⁹⁷ protein a T cell proliferative assay was performed as previously described by Bixler and Atassi (Immunol. Commun. 12:593, 1983). Briefly, SJL/J mice were immunized with 50 µg of native CRM¹⁹⁷ emulsified in CFA. Seven days later, lymph nodes were removed, cultured in RPMI and challenged with various concentrations of proteins (0.05-100.0 µg/ml) and peptides. After 3 days incubation, cultures were pulsed with [³H]-thymidine for 16 hours and then harvested for counting.

TABLE 3

T cell responses to meningococcal peptide-CRM197 conjugates.
In Vitro Challenge

Maximum observed [^3H] incorporation	$\mu\text{g/ml}$	dCPM	SI
<hr/>			
Diphtheria toxoid CRM197 - mock conjugate	5	27,510	57
	50	108,631	221
	100	116,326	236
	100	182,499	370
	10	89,972	183
M21-CRM197	100		
M20-CRM197	10		
CON A LPS Tetanus toxoid	1	34,316	70
	50	61,579	126
	10	515	2
Background (cpm)	-	494	1

As shown in Table 3, a comparison of CRM₁₉₇ with the CRM₁₉₇-mock conjugate shows that the conjugation procedure by itself did not alter the T cell recognition of the protein. The T cell responses induced by the M20 and M21-CRM₁₉₇ conjugates were essential equivalent to or greater than the response elicited by CRM₁₉₇ itself indicating that the recognition of the T cell epitopes on the CRM₁₉₇ is not adversely affected by the peptide conjugation. The responses to the control materials Con A, LPS and Tetanus toxoid were as expected.

EXAMPLE 13: Immunogenicity of conjugate and recombinant meningococcal vaccines

Recombinant flagellin expressing the meningococcal VR1 and/or VR2 epitopes were prepared and purified as described in Examples 7, 8 and 9. In addition, synthetic peptides representing the meningococcal epitopes VR1 and VR2 were synthesized, covalently coupled to the carrier molecule CRM₁₉₇ and purified as in Example 12. Vaccines were formulated with each of these materials at protein concentrations of 10 or 100 µg/ml for each of the components. The vaccine compositions also included aluminum phosphate at 1 mg/ml or except as noted were compounded with Freund's complete adjuvant or without supplemental material. To evaluate immunogenicity, outbred Swiss Webster mice were immunized intramuscularly at weeks

0 and 2 with 1 or 10 μ g protein/dose. Sera were collected at two week intervals, pooled for assay, and screened for antibody activity by ELISA to outer membrane complex (OMC), purified OMP (PI.16), VR1 peptide coupled to Bovine serum albumin (M21-BSA), VR2 peptide coupled to BSA (M20-BSA), wildtype flagellin, and to CRM₁₉₇. The results of the ELISA performed on sera obtained at 6 weeks are shown in Table 4.

DOSE	#9	OMC	P1.16	M21-BSA	M20-BSA	FLAGELLIN	CRM
ELISA TITERS 4 WEEKS AFTER SECONDARY BOOST-							

PCB1-4	PCB2-W
10	10
1	1
532	150
2,034	1,350
4,376	308
12,387	12,190
4,525	ND
17,565	ND
ND	5,476
ND	501
787,120	263,143
887,861	1,493,216
ND	ND

Category	Sub-category	Value	Unit
M20-CRM197	10	150	<100
M20-CRM197	1	217	<100
M20-CRM197	10	50	<100
M21-CRM197	1	249	311
M21-CRM197	10	10,494	26,807
M21-CRM197	1	100	191
M21-CRM197	10	17.41	20.92
M20-CRM197	1	42.27	95.31
M20-CRM197	10	ND	ND

PCB1-4 in CFA				PCB2-W in CFA			
10	1,665	10,606	19,945	ND	17,749	1,217,063	ND
1 All pre-bleed values at or below the lower limit of assay of 1							

2 All vaccines were formulated with 1 mg/ml aluminum phosphate solution. as not d.

-71-

Alternatively, the various vaccines were evaluated for immunogenicity in 6-8 week old NIH outbred mice. The mice were immunized with 100 µg (total protein)/dose subcutaneously on week 0 and 4 with vaccine and sera was collected on week 6. The sera were evaluated in an ELISA assay and using antigens as described in Example 6. Bactericidal activity was measured as in Example 6. The results are found in Table 5.

TABLE 5
ELISA (titer > 0.5 OD)

Bactericidal Test	Synth. Peptide	Class 1 OMP			OMC			Vaccine	
								PLAGELLIN	p1650
<1:64	-	-	-	-	-	-	-	pCB12.10.6	pCB2-V
<1:64	1:100	1:300	1:2700	1:300 (.25)	-	-	-	PCB1-4	CRM197
<1:64	-	-	-	-	-	-	-	H20-CRM197	H21-CRM197
<1:64	-	-	-	-	-	-	-		
<1:64	-	1:8100	1:8100	1:300 (.125)	1:00	1:300	1:125		

The recombinant flagellins containing either a

VR1, VR2 or a cassette of both VR1 and VR2 were

effective in eliciting an antibody response which

was cross-reactive to the purified P1.16 and to a

lesser extent to OMC. Sera from animals immunized

with 10 µg of either PCB1-4 or PCB2-W induced

antibodies which bound to their respective peptide-

-BSA conjugates as well as cross reacted with the

P1.16 and OMC. Similar results were obtained with

the constructed PCB12-10-6 which contains both

meningococcal epitopes. In addition, each cons-

truction induced significant anti-flagellin titers

as well. In contrast, the control wildtype fla-

gellin only induced an antibody response to fla-

gellin itself. Sera collected prior to immunization

showed no pre-existing response to the materials

being evaluated.

The data also demonstrates the benefits of

formulating the recombinant flagellins with alum or

other adjuvants such as CFA. The construction

PCB12-10-6 was formulated with and without the

addition of aluminum phosphate. As shown in table

2, PCB12-10-6 alone was capable of inducing an

antibody response which react to the peptide con-

jugates as well as to the purified P1.16 as well as

to OMC. In comparison, the same construction when

formulated with alum was able to elicit greater

antibody response at an equivalent dose. Similarly,

the recombinant flagellins PCB1-4 and PCB2-W were

also formulated with CFA. Again, equivalent or

-73-

higher antibody titers were observed in the presence of CFA.

The results of the immunogenicity studies with the meningococcal VR1 and VR2 conjugates are also shown in Table 4. Both the M20 and the M21-CRM¹⁹⁷ conjugates as well as a mixture containing equal amounts of both conjugates were capable of inducing an anti-CRM¹⁹⁷ response as well as an anti-Class I OMP response.

These preliminary data indicate a Class I OMP

variable region epitopes either chemically

conjugated to a carrier or genetically fused to a

carrier elicit an immune response. New epitope-

carrier conjugates can be made using standard

techniques to enhance the immune response to the

vaccine, for example, use of 1) larger epitopes, 2)

peptides with multiple epitope repeats and/or 3)

different carriers.

EXAMPLE 14: Preparation of Meningococcal-human serum

albumin-lycocalmucate

GCM CPS was depolymerized by acid hydrolysis and GCM OS obtained were subsequently activated via NaIO_4 oxidation in aqueous buffer. The reactions were monitored by HPLC in aqueous eluent, using UV and RI detection. The reactions were each followed by GPC desalting in water. GCM OS and human albumin (HA) were mixed and conjugated essentially as described in Example 10 for the meningococcal-

flagellin glycoconjugate. The final preparation was stored in the cold, in the presence of thimerosal to prevent bacterial growth.

In preparation of the conjugate, the following experimental conditions were employed. Human

albumin (HA; Sigma^R, St. Louis, MO) was dissolved in 15% sucrose (10 mg/ml) and then stored at -20°C.

GCM CPS (lot # 86 NM 01; 106 mg; final concentration: 10 mg/ml was hydrolyzed in 0.1 N HCl at

50°C with agitation. Aliquots (25 μ l) were withdrawn at regular intervals, the reaction stopped by

addition of sodium hydroxide (NaOH) and analysis was performed by HPGPC as described. After 3 h 40 min.,

the reaction was stopped by addition of NaOH, and the GCM OS were desalted by GPC. Fractions were

collected (1.2 ml) and analyzed as described before. Positive fractions were pooled and lyophilized.

Desalted GCM OS (89 mg) were then stored at -20°C. Activated OS were prepared by oxidation of GCM OS

(11.8 mg; final concentration: 5mg/ml) with 2 mM NaIO₄ in 0.05 M sodium phosphate buffer (pH 6.2-6.5)

at RT, in the dark, with agitation. The reaction was stopped after 30 min by addition of ethylene

glycol. HPGPC analyzes showed no degradation of the molecular weight of the OS during activation.

Desalting and colorimetric analyzes were then performed as described above. The resulting

activated GCM OS (8.8 mg) were dissolved in water (10 mg/ml) and frozen at -20°C.

-75-

Both GCM OS and HA solutions were analyzed by HPLC (UV at 206 and 280 nm respectively) before being frozen, and prior to the conjugation. No degradation occurred during storage, as ascertained by the exact similarity of the elution profiles. GCM OS (6 mg; final concentration: 2.5 mg/ml) and HA (12 mg; final concentration: 5 mg/ml) were mixed in a polypropylene tube in 0.4 M sodium phosphate buffer (pH 7.0), and NaBH_3CN was added (60 μmoles) to initiate the conjugation (Anderson, U.S. Patent 4,762,713, 1988; U.S. Patent 4,673,574, 1987; U.S. Patent 4,761,283, 1988). The reaction mixture was left one day at RT, then 4 days 15 35°C , without agitation. The reaction was monitored by HPLC (UV at 280 nm) at different stages, and finally stopped by dialysis/concentration on microconcentrators. The final preparation was analyzed for NANA (Barry et al., J. Gen. Microbiol., 29P335-51, 1962) (2.07 mg at 0.86 mg/ml) and protein (Lowry et al., J. Biol. Chem., 193265-75, 1951) (9.51 mg at 3.96 mg/ml) content. It was then stored at 4°C in the presence of thimerosal (0.01%, w/v) to prevent bacterial growth. The conjugate preparation was also checked by SDS-PAGE (silver nitrate stain) and Western blot analyzes. A diffuse band appeared on the gel which covered a significantly wider molecular weight range than the pure HA. Western blot analyzes showed that this band was reactive with the antiserum used

(anti-GCM), proving a valency of the conjugate bonds.

EXAMPLE 15: Immunogenicity of meningococcal

oligosaccharide-recombinant flagellin

vaccines

A meningococcal oligosaccharide-recombinant flagellin vaccine was prepared as described above and formulated at 100 µg protein/ml. Vaccine compositions were also prepared which contained aluminum phosphate (1 mg/ml) or complete Freund's adjuvant in addition to the glycoconjugate.

To evaluate the immunogenicity, outbred Swiss Webster mice were immunized intramuscularly with 10 µg protein at week 0 and 2. Sera were collected at weeks 0, 2 and then weekly intervals thereafter to 6 weeks. After collection, pooled sera samples were assayed for antibody activity by ELISA to meningococcal C oligosaccharide conjugate to human serum albumin, OMC, PI.16, CBI and CB2-BSA conjugates and flagellin.

The MENC-CBI2-10-6 glycoconjugate was effective at eliciting an immune response which was reactive with both the oligosaccharide and the meningococcal B OMP epitopes expressed in the recombinant flagellin. As shown in Table 5B, as little as three weeks into the study, mice immunized with 1 µg of MENC-CBI2-10-6 conjugate in complete Freund's adjuvant had detectable antibody to MENC-HSA, OMP

and to both the CB1 and CB2 epitopes. Further, all of the MENC-CB12-10-6 preparations, regardless of adjuvant, elicited antibody response to MENC-HSA which were greater than the response observed following immunization with MENC-CRM197.

Table 5B. Immunogenicity of Meningococcal C recombinant flagellin vaccine one week after secondary immunization.

IMMUNOGEN									
Dose MENC-HSA									
ELISA TITERS									
OMP CB1-BSA CB2-BSA FLAGELLIN									
MENC-CB12-10-6	CFA	10	24,530	608	5,240	5,614	5,375	12,685	432
		1	5,069					526,593	541,467
	alum	2	11,845	253	835			673	472,766
		1	4,415	136	242			244	214,263
	None	10	11,497	920	626			2,382	233,307
		1	4,920	483	1,123			1,210	135,625
MENC-CRM197	alum	10	4,905	ND	ND	ND	ND	ND	ND
		1	8,505	ND	ND	ND	ND	ND	ND
OMP (PI.16)	alum	10	ND	12,907	<100	<100	<100	3,377	ND
		1	ND	10,405	<100	<100	<100	ND	ND

1 Titers for initial prebleed samples (week 0) samples were <100.
2 Aluminum phosphate was used as adjuvant at 1 mg/ml.

EXAMPLE 16: T-cell epitopes of Class I OMP and their identification

An effective vaccine must contain one or more

T-cell epitopes. T-cell epitopes within a protein can be predicted as described by Margalit *et al.*, J. Immunol. 138:2213, (1987) or Rothbard and Taylor, EMBO J. 7:93, (1988). These predictive methods were

applied to the amino acid sequence of the Class I OMP of *N. meningitidis* strains PI.7, PI.16 and PI.15. The segments of the sequence containing

potential T cell epitopes identified by these methods are shown in Tables 6 and 7. The predicted

peptides were synthesized by standard Fmoc

procedures, purified by standard methods and were identified as shown in Table 8.

To determine which of the predicted peptides

actual contain T cell epitopes, their capacity to

stimulate human peripheral blood lymphocytes (PBL)

was tested by lymphocyte proliferative assay.

Briefly, peripheral blood was collected from HLA

typed normal volunteers or from volunteers who were

previously immunized with MPC-2 (Poolman, J.T. *et*

al., *Antonie van Leeuwenhoek*, 53:413-419, 1987)

which contained PI.16, PI.15, Class 4 OMP and Group C

polysaccharide. Lymphocytes were isolated from the

peripheral blood by isolation on Ficol-Hypaque

(Pharmacia Fine Chemicals AB, Uppsala, Sweden) and

cultured at 1×10^5 cells/well in supplemented RPMI

1640 (Gibco Laboratories, Paisly, Scotland)

containing 10% heat-inactivated pooled human AB serum. Cultures were challenged with various concentrations of the predicted T cell epitopes (0.05 - 10 μ g/ml). After *in vitro* challenge, the cultures were incubated for six days and then pulsed (18 hours) with 0.5 μ ci of [3 H]-thymidine. Cultures were then harvested and counted by liquid scintillation. Data are expressed as stimulation indices which were calculated as a ratio of the CPM obtained in the presence of antigen to the CPM obtained in the absence of antigen.

As shown in Table 9, 10 of the 16 predicted peptides showed some capacity to stimulate T-cells. These include the peptides identified at 16-34, 47-59, 78-90, 103-121, 124-137, 151-158, 176-185, 223-245, 276-291 and 304-322. In some instance, peptides stimulated a response in both immunized as well as non-immune individuals. The response in the non-immune individuals may be attributed to a previous asymptomatic infection.

In the case of the T cell epitope identified as region 176-185, enhancement of the T cell response was observed following addition of the monoclonal antibody MN5C11G (P1.16). Briefly, PBL were challenged *in vitro* with a synthetic peptide containing the region 175-185 or with this peptide mixed with varying dilutions of MN5C11G. As shown in Table 10, enhancement of the T cell response was observed following addition of MN5C11G indicating that monoclonal antibody recognized a B cell epitope

within the region 176-185 and facilitates the presentation of the peptide to the immune system. Thus, it was established that the T and B cell epitopes either coincide or are found on contiguous sequences within the Class I OMP.

In several cases, T cell lines and clones were established from individuals responding to various peptides. Briefly, T cell lines were obtained by culturing isolated lymphocytes in 24 well plates at 1×10^6 cells/ml. The culture medium, supplemented RPMI-1640 with 10% human serum, also contained 12 U/ml recombinant IL-2 (Boehringer). In addition, 5×10^4 homologous, irradiated ($3,000R$) antigen presenting cells (APC) were also added to each well. In some cases, APC were obtained from HLA compatible donors. From the lines, T cell clones were isolated by limiting dilution at a frequency of 0.5 cells/well. Clones were maintained by bi-weekly stimulation with antigen in the presence of irradiated APC and IL-2 (2 U/ml). Clones were tested by lymphocyte proliferation assay essentially as described above except that clones were cultured at 1×10^4 cells/well in the presence of irradiated APC.

Clones obtained as described were challenged in vitro with OMP from 7 different strains of meningococci. As shown in Table II, the clones recognized a T cell epitope or epitopes common to the seven OMPs examined. Although the reactivity of these clones to the various peptides remains to be

Table 6. ANALYSIS OF THE SEQUENCE OF N. MENINGITIDIS P1.16 OMP
FOR THE PRESENCE OF AMPHIPATHIC α -HELICES ACCORDING
TO THE METHOD OF MARGALIT ET AL. (J. IMMUNOL. 136:2213, 1987)

MID POINTS OF BLOCKS	ANGLES	AS
47-50	85-105	9.4
69-74	105-135	16.0
79-88	90-120	23.0
127-135	100-120	22.4
199-202	90-120	8.4
208-211	85-95	8.7
260-263	90-125	8.8
265-269	90-120	11.3
274-277	105-120	9.8
297-300	100-135	9.1
320-324	80-100	10.9
338-342	105-135	12.3
346-351	80-115	11.9
376-379	85-120	9.5

N R K K L T A L S A L P L A V A D V S L G E I K A G V E G R N I
 Q A O L T E Q P O V T N G V G N Q V K V I K A K S R I K I S D I G
 S F I G F K G S E D L G E G L K A V M Q L E Q D V S V A G G A S Q M G
 N R E S F I G L A G E F G T L R A G R V A N Q F D D A S Q A I N P M D S
 N N D V A S Q L G I F K R H D P V S V K Y D S P F S G F S G S V Q
 P V P A Q N S K S A Y K P A Y T K D T N N N L T L V P A V G K P G S
 D V Y A G L N Y K N G G P A G N Y A F K X A R A N V G R N A F E L F
 L I G S A T S D E A K G T D P L K N H Q V H R L T G G Y E E G G L N L A
 L A V Q L D L S E N G D K A K T K N S T I E I A V T A S Y R I G N A V P
 M I S Y A H G F D L I E R K G K G E N T S Y D Q I I A G V D X D I S K R
 T S A I V S G A M L K R N T G I G N Y T Q I N V A S V G L R E K F

Table 7. PRESENCE OF MOTIFS (UNDERLINED REGIONS) REPRESENTING
 POTENTIAL T CELL EPITOPES WITHIN THE SEQUENCES OF N. MENINGITIDIS
 P1.16 OMP AS DETECTED BY THE METHOD OF
 ROTHBARD AND TAYLOR (EMBO J 7:93, 1988).

Table 8. SUMMARY OF PREDICTED T CELL EPITOPES SYNTHESIZED.

SEQUENCE		RESIDUE NO.
<hr/>		
NIQALTEQPVNTNGVGN	16-34	1.
TKISDFGSFIGFK	47-59	2.
GFKGSEDLGEGLKAV	57-71	3.
VSVAAGGASQWGN	78-90	4.
TLRAGRVAHQFDDASQAIN	103-121	5.
DSNNDVASQLGIFK	124-137	6.
GGFSGFSG	151-158	7.
YTTKDTNNNL	176-185	8.
AVVGKPGSDVYYA	190-202	9.
YAFKYYARNAHVGRN	215-228	10.
ANVGARNAFELFLIGSATSDKAG	223-245	11.
DEAKGTDFLKNHQVHRLTGGY	241-261	12.
LSENGDKAKTKNSTTE	276-291	13.
VPRISSYAHGFDLIERGKKG	304-322	14.
ERGKKGENTSYDQ	317-329	15.
KRNTGIGNYTIQINAA	352-366	16.

Table 9. SUMMARY OF LYMPHOCYTE RESPONSES TO MENINGOCOCCAL SYNTHETIC PEPTIDES IN HLA TYPED VOLUNTEERS.

RESPONSE TO SYNTHETIC PEPTIDE															
VOLUNTEER/HLA TYPE															
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
IMMUNIZED VOLUNTEERS															
1. DR4, W10, W53	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2. DR3, W52	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
3. DR3, W52, W53	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4. DR2, W6, W13, W15, W52	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5. DR5, W11, W7, W52, W53 +	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
6. DR5, W11, W10, W52	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
NON-IMMUNIZED CONTROL VOLUNTEERS															
7. NOT TYPED	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
8. DRW13, W6, W52	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
9. DR2, W15, W4, W53	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10. NOT TYPED	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
11. NOT TYPED	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
12. DR2, W15, W3, W52	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
13. DR5, W11, W52	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
14. DR3, W7, W52, W53	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15. DR3, W4, W52, W53	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
16. DR3, W12, W5, W52	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
17. DR2, W15, W7, W53	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
18. DR1, W3, W52	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
19. DR3, W4, W52, W53	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20. DR1, W7, W53	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
21. DR4, W8, W52, W53	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
22. DR1, W13, W6, W52	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
23. DR2, W16, W5, W11, W52	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
24. DR5, W11, W6, W13, W52	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
25. DR1, W3, W52	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
26. DR1, W6, W13, W52	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
27. DRW6, W13, W52	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
28. DRW6, W13, W52	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

The responses were scored as follows - , SI<2; +, 2<SI<3 and +, SI > 3.

Table 10. PRESENTATION OF A SYNTHETIC PEPTIDE TO PERIPHERAL BLOOD LYMPHOCYTES IS ENHANCED BY A MONOCLONAL ANTIBODY RECOGNIZING REGION 179-184 OF MENINGOCOCCAL CLASS I OMP.

IN VITRO CHALLENGE		CPM
GGYYTKDITNNNL		3,017
GGYYTKDITNNNL* + MNSC11G (1:200)		22,836
GGYYTKDITNNNL + MNSC11G (1:1000)		12,608
MEDIA		330

*Underline region indicates sequence recognized by monoclonal antibody MNSC11G.

STRAIN SUBTYPE	5-5	5-7	5-9	5-12	5-13	5-14	5-15
H44-76	P1.16	6.0	1.2	6.8	2.6	2.3	9.5
SWISS 4	P1.15	4.9	1.0	10.1	6.9	3.6	10.5
395	P1.9	5.2	1.5	4.8	1.5	6.1	13.1
2996	P1.2	5.4	1.0	3.7	2.3	3.4	11.8
H990	P1.6	3.6	0.4	3.5	2.5	0.9	4.7
187	P1.1	4.4	0.7	4.5	3.1	1.6	6.2
6557	P1.17	3.7	2.0	8.2	4.2	1.7	6.2
MEDIA	--	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1

Table 11. RECOGNITION OF OMP FROM DIFFERENT MENINGOCOCCAL STRAINS BY HUMAN T-CELL CLONES

EXAMPLE 17:
Construction of Protein Model for
Membrane Topology of Class I OMP and
Comparison to Other Pathogenic Gram
Negative Porin Proteins for Vaccine
Development.

A model was constructed using the principles recognized for the structure of several Escherichia coli outer membrane proteins (Vogel, H. et al. (1986) supra Ferenci, T. et al. (1988) supra; and Tommassen, J. (1988) supra). The central assumption is that protein segments spanning the outer membrane form beta-sheets. Specifically, in the case of Class I protein, the division in exposed and trans-membrane segments was arrived at in the following way:

1. A comparison of the amino acid sequence of Class I protein (subtype PI.16) with those of the gonococcal PIA and PIB proteins (Carbon-etti, N.H. et al. (1987) PNAS 84:9084; Carbon-etti, N.H. et al. (1988) PNAS 85:6841; and Gotschlich, E.C. et al. (1987) PNAS 84:8135) reveals 34% identity. In the model, the variable sequences form the surface-exposed parts, whereas the conserved regions are placed mostly in the outer membrane and periplasm. Thus, the latter two areas consist for 58% of residues that are conserved among all proteins.

2. The hydrophilic maxima observed in a hydrophathy profile (Kyte, J. et al. (1982) J. Mol. Biol. 157:105) to correspond to exposed regions.

3. The transmembrane segments should preferably be able to form amphipathic beta-stands of 9-12 residues, with at least one side consisting entirely of hydrophobic residues. These conditions are met in 12 of the 16 membrane-spanning segments.

4. The number of residues at the periplasmic side is minimized.

Figure 11 shows the model for the folding of Class I protein in the outer membrane. The sequence shown is for subtype Pl.16. The top part of the figure shows the surface-exposed regions, whereas the central part indicates the presumed trans-membrane segments, whose length is set at ten. Amino acid are shown alternating where they can form an amphipathic beta-strand. This model contains eight surface loops, whereby the first and the fourth loop contain the type-specific and protective epitopes. These epitopes, as has been shown when formulated into a vaccine, can elicit a protective immune response. Loop 5 is constant and has been shown to elicit cross-reactive antibodies to other OMPs and is useful for vaccine formulation.

The one or two variable epitope regions of the individual proteins are located on so called surface loops of these membrane proteins. Such porin outer-membrane proteins contain more than two surface loops. This implicates that there are surface loops which have near identical amino acid sequence in the different Class I outer-membrane proteins as well. This opens the way to use of common peptides of the Class I outer-membrane protein for vaccine objectives as well. More especially a schematic two-dimensional model of the meningococcal Class I outer-membrane protein P1,16 is illustrated in Figure 11. This model contains eight surface loops, whereby the first and the fourth loop contain the type specific epitopes as shown on the basis of strain subtyping results. The fifth surface loop represents the constant region described above. Antibody to the constant region of loop 5 appears to react with *N. meningitidis* OMP complex. The amino sequence of Class I OMP, as derived, was compared to the Class OMP of *N. meningitidis* (Mura-kami, K. et al., (1989), *Infect. Immun.*, 57:2318) and the porin P1A and P1B proteins of *N. gonorrhoeae*. With similar principle as used for the Class I OMP modeling, the sequences were aligned as follows:

LOOP 1

Class I DVLGYEIKAG VEGRNIOAQLTEQVLTNGVQGN

Class IB DVLGYAIIKAG V-----QTYRSVEHT

Class IA DVLGYTIKAG VEGRNIOAQLTEQFETSRVAHH

Class II DVLGYTIKAG EGRNIOAQLTEQFEVSRVKDAG

QVKVTKAKSRIRTKI SPFGSTIGFKGSEDLGEGT

DGVSKVET--GSEI ADFGSKIGFKGQEDLGNGL

GAQADRVT--ATEI ADLGSKIGFKGQEDLGNGL

TYKAQGGKSKTATQ IADFGSKIGFKGQEDLGNGL

LOOP 2

KAVWOLEQD VSAAGGASQWGN RESFGLAGEFG

KAVWOLEQD ASVAGTNTG-WGN KQSFVGLKGGFG

KAVWOLEQD AYVSGTDTG-WGN RQSFGLKGGFG

KAIWOLEQ KASIAGTNSG-WGNRQSFGLKGGFG

LOOP 3

TLRAGRVAHQFDD ASQAINFWDNNDDVASQLOI-

TIRAGSLNSPLKN TGANVANAWESGKFTGNVLZIS

KVRVGRHTNSYTKD TGGFNPFWEKSYLPLTNSIAQ

TVKAGNTLNTVLKDSGDNNANAWESGSNTEDVLGLG

RVAVNTANAHFVKDY *****
* OVRVAVAGYDNDLYSVA *

FSKRISALVSAGWL KOGKGTGKVEQ-LASM VGLRHKF *****
* * * * *

FSKRISALVSAGWL KOGKGTGKVEQ-LASM VGLRHKF ***** * * * * *	FSKRISALVSAGWL KOGKGTGKVEQ-LASM VGLRHKF ***** * * * * *
FSKRISALVSAGWL KOGKGTGKVEQ-LASM VGLRHKF ***** * * * * *	FSKRISALVSAGWL KOGKGTGKVEQ-LASM VGLRHKF ***** * * * * *
FSKRISALVSAGWL KOGKGTGKVEQ-LASM VGLRHKF ***** * * * * *	FSKRISALVSAGWL KOGKGTGKVEQ-LASM VGLRHKF ***** * * * * *
FSKRISALVSAGWL KOGKGTGKVEQ-LASM VGLRHKF ***** * * * * *	FSKRISALVSAGWL KOGKGTGKVEQ-LASM VGLRHKF ***** * * * * *

LOOP 6

TPRVSYAHEF KANVNGVKDANYQDDQ VIAGADYD *****
* * * * *

TPRVSYAHEF KANVNGVKDANYQDDQ VIAGADYD ***** * * * * *	TPRVSYAHEF KANVNGVKDANYQDDQ VIAGADYD ***** * * * * *
TPRVSYAHEF KANVNGVKDANYQDDQ VIAGADYD ***** * * * * *	TPRVSYAHEF KANVNGVKDANYQDDQ VIAGADYD ***** * * * * *
TPRVSYAHEF KANVNGVKDANYQDDQ VIAGADYD ***** * * * * *	TPRVSYAHEF KANVNGVKDANYQDDQ VIAGADYD ***** * * * * *
TPRVSYAHEF KANVNGVKDANYQDDQ VIAGADYD ***** * * * * *	TPRVSYAHEF KANVNGVKDANYQDDQ VIAGADYD ***** * * * * *

LOOP 7

GQ YEAKNNVEGSTRGKKHQTO VAAATAYRFGNV *****
* * * * *

GQ YEAKNNVEGSTRGKKHQTO VAAATAYRFGNV ***** * * * * *	GQ YEAKNNVEGSTRGKKHQTO VAAATAYRFGNV ***** * * * * *
GQ YEAKNNVEGSTRGKKHQTO VAAATAYRFGNV ***** * * * * *	GQ YEAKNNVEGSTRGKKHQTO VAAATAYRFGNV ***** * * * * *
GQ YEAKNNVEGSTRGKKHQTO VAAATAYRFGNV ***** * * * * *	GQ YEAKNNVEGSTRGKKHQTO VAAATAYRFGNV ***** * * * * *
GQ YEAKNNVEGSTRGKKHQTO VAAATAYRFGNV ***** * * * * *	GQ YEAKNNVEGSTRGKKHQTO VAAATAYRFGNV ***** * * * * *

LOOP 6

- 93 -

Structural similarities are indicated with transmembrane and surface loop regions. With the information now available for Class I OMP and intraspecies amino acid homology or heterology of the loop regions of the particular porin protein, predictions of epitopes for incorporation into vaccines for other pathogenic gram negative bacteria including *N. gonorrhoeae* are possible. Using the same methods employed for Class I OMP, these epitopes can be evaluated for vaccine purposes.

Deposit of Microorganisms

The *N. meningitidis* strain H4476 (B:15:P1.7,16) was deposited on December 11, 1989 in the Central Bureau voor Schimmelculturen (CBS), Baarn, The Netherlands and has deposit number CBS 635.89. The *N. meningitidis* Class 2/3 OMP deficient mutant H111-5 was deposited on December 11, 1989 in the CBS, Baarn, The Netherlands and has deposit number CBS 636.89.

Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

CLAIMS

1. Vaccine effective against meningococcal disease, comprising an effective amount of an outer-membrane vesicle isolated from expressing homologous and/or heterologous or at least one meningococcal class I outer-membrane protein or a fragment or oligopeptide containing an epitope thereof.
2. A vaccine of Claim 1, wherein the meningococcal class I outer-membrane protein originates from a mutant meningococcal strain which is negative for Class 2/3 outer membrane protein.
3. A vaccine of Claim 1, wherein the Class I outer-membrane protein, fragment or oligopeptide is produced by a microorganism containing a heterologous gene encoding the Class I outer-membrane protein, fragment or oligopeptide.
4. A vaccine of Claim 1, wherein the Class I outer-membrane protein, fragment or oligopeptide is derived from a group A, B, C, W-135 or Y meningococcus.
5. A vaccine of Claims 1, wherein the Class I outer-membrane protein, fragment or

oligopeptide is derived from a group B meningococcus.

6. A vaccine of Claim 1, comprising 5-10 various group B meningococcal class I outer-membrane proteins, fragments or oligopeptides.

7. A vaccine of Claim 1, wherein the protein fragment is obtained by cyanogen bromide treatment of a Class I outer-membrane protein.

8. A vaccine of Claim 1, wherein the fragment of class I meningococcal outer-membrane protein is obtained by proteolysis with an enzyme selected from the group consisting of endolysins, endo-Arg-C, endoglycosidase and Staphylococcus V8-protease.

9. A vaccine of Claim 1, wherein the oligopeptide comprises at least one bactericidal antibody binding epitope of meningococcal class I outer-membrane proteins.

10. A vaccine of Claim 1, wherein the oligopeptide comprises at least an amino acid sequence selected from the group consisting of QPQVTNGVQGN, PPSKSP, QAAANGGASG, YTKDITNNLTL, YTKNTNNLTL, YTKDITNNL, YTKNTNNL, HFVQQTTPQSP and HYTRQNNNTDVF.

11. A vaccine of Claims 1, wherein the epitope is located in surface loops of meningococcal class I outer-membrane proteins in the area of amino acids 24-34 and 176-187.
12. A vaccine of Claim 1, wherein the Class I outer-membrane protein, fragment or oligopeptide is conjugated to a T cell epitope via chemical coupling.
13. A vaccine of Claim 1, wherein the Class I outer-membrane protein, fragment or oligopeptide contains a meningococcal A,B,C,W-135 and/or Y polysaccharide in conjugate form with the protein product.
14. A vaccine of Claim 1, further comprising the a Zwitterionogenic, cationogenic, anionogenic and/or non-ionogenic detergent.
15. A vaccine of Claim 14, wherein the detergent is selected from the group consisting of Zwittergent ZW 3-10, Zwittergent ZW 3-14, Tween-20, sodium cholate, octyl glucoside and sodium deoxycholate.
16. A vaccine of Claim 1, further comprising the an adsorbent selected from group consisting of aluminum phosphate, aluminum hydroxide and calcium phosphate.

23. Oligopeptide containing a B cell epitope of a

22. A fragment of Claim 21, produced by cyanogen bromide cleavage of a class I outer membrane protein of *N. meningitidis*, the fragment having a molecular weight of approximately 25 kD.

21. A fragment of Claim 20, wherein the class I outer membrane protein is of the subtype Pl.7.16.

20. Substantially purified fragment of class I outer membrane protein of *Neisseria meningitidis*, the fragment having a molecular weight of about 25 kD or less and containing continuous or discontinuous epitopes reactive with bactericidal antibodies against *N. meningitidis*.

19. A vaccine of Claims 1, the Class I outer membrane protein, fragment or oligopeptide is coupled to a lipid.

18. A vaccine of Claim 1, wherein the Class I outer membrane protein, fragment or oligopeptide is contained within a liposome.

17. A vaccine of Claim 1, further comprising an immuno-stimulating complex (ISCOM).

28. A method of eliciting a protective immune response against Neisseria meningitidis, comprising administering a vaccine composition, comprising one or more meningococcal Class I outer membrane proteins or fragments thereof and, optionally, an adjuvant in a pharmaceutically acceptable vehicle.

27. Isolated nucleic acid encoding meningococcal Class I outer-membrane protein or a fragment or oligonucleotide containing an epitope thereof.

26. Oligopeptide containing a T cell epitope of a meningococcal Class I outer-membrane protein.

25. Oligopeptide containing an epitope of a Class I outer-membrane protein which is conserved in amino acid sequence among different meningococcal porin proteins.

24. Oligopeptide of Claim 23, containing at least one of the amino acid sequences selected from the group consisting of PQVTNGVQGN, PPSKSP, QAANGGASG, YTKDNTNNLTL, YTKNTNNLTL, YTKDNTNNL, YTKNTNNL, HFVQQTSPQSP and/or HYTRQNNITDVF.

Class I outer-membrane protein which varies in amino acid sequence among different meningococcal porin proteins.

29. A method of Claim 28, wherein the fragment is conjugated or genetically fused to a T cell epitope, B cell epitope or carrier peptide or protein.
30. A method of Claim 29, wherein the conjugation is through a cysteine or lysine residue coupled to a terminus of the fragment.
31. A method of Claim 29, wherein the carrier is a bacterial toxin, CRM or toxoid.
32. A method of Claim 28, wherein the class I outer membrane protein is of subtype PI.7,16.
33. A method of eliciting a protective immune response against *Neisseria meningitidis*, comprising administering a vaccine composition, comprising at least one oligopeptide containing a continuous or discontinuous epitope of a class I outer membrane protein reactive with bactericidal antibodies against *N. meningitidis* and, optionally, an adjuvant in a pharmaceutically acceptable vehicle.
34. A method of Claim 33, wherein the oligopeptide contains an amino acid sequence selected from the group consisting of QPQVTNGVQGN, PPSKSP, QAANGGASG, YYTKDITNNLTL, YYTKNTNNLTL, YYTKDITNNNL, YYTKNTNNNL, HFVQQTTPQSP and/or HYTRQNNITDVF.

35. A method of Claim 33, wherein the proteolytic fragment is conjugated or genetically fused to a T cell epitope, B cell epitope or carrier peptide or protein.
36. A method of Claim 33, wherein the conjugation is through a cysteine or lysine residue coupled to a terminus of the fragment.
37. A method of Claim 35, wherein the carrier is a bacterial toxin, CRM or toxoid.
38. An antigenic conjugate, comprising a meningococcal class I outer-membrane protein, a fragment or oligopeptide containing an epitope thereof conjugated to a carrier protein or epitope thereof.
39. An antigenic conjugate of Claim 8, wherein the oligopeptide selected from the group consisting of QPQVINGVQGN, PPSKSP, QANAGASC, YTKDITNNLTL, YTKNITNNLTL, YTKDITNNL, YTKNITNNL, HGVQQTTPQSP and/or HYTRQNNITDVF.
40. An antigenic conjugate of Claim 39, wherein the antigen carrier protein is a bacterial toxin, CRM or epitope thereof.
41. An antigenic conjugate of Claim 40, wherein the carrier protein is CRM¹⁹⁷.

42. A genetic fusion peptide or protein, comprising an epitope of an meningococcal class I outer-membrane protein fused to a carrier protein, peptide or epitope thereof.

43. A genetic fusion peptide or protein of claim 42, wherein the epitope is selected from the group consisting of QPQVTNGVQGN, PPSKSP, QVANGASG, YTKDITNNLTL, YTKNTNNLTL, YTKDITNNNL, HFTVQQTPOSP and/or HYTRQNNITDVF.

44. A fusion protein, comprising a flagellin protein having an amino acid sequence for an epitope of a meningococcal class I outer-membrane protein inserted within it.

45. The fusion protein of claim 44, wherein the amino acid sequence is inserted within the flagellin protein at a region which is non-essential to function of the flagellin protein.

46. The fusion protein of claim 45, wherein the amino acid sequence is inserted into the hyper-variable region of the flagellin protein.

-102-

47. A fusion protein of Claim 44, further comprising a meningococcal capsular oligo- or polysaccharide conjugated thereto.
48. A recombinant gene, encoding a fusion protein of Claims 44.
49. An infectious, recombinant microorganism capable of expressing a class I outer membrane protein of Neisseria meningitidis, or a fragment or oligopeptide containing a continuous or discontinuous epitope reactive with bactericidal antibodies against N. meningitidis.
50. A microorganism of Claim 49, wherein the oligopeptide selected from the group consisting of QPQVINGVQGN, PPSKSP, QANAGASC, YTKDNTNNLT, YTKNTNNLT, YTKDNTNNLT, YTKNTNNLT, HFVQQTTPQSP and/or HYTRQNNTDVF.
51. A microorganism of Claim 49, which is a vaccinia virus, adenovirus, or cytomegalovirus.
52. A microorganism of Claim 49, which is a bacteria of the genus Salmonella.
53. A microorganism of Claim 49, wherein the epitope of the meningococcal class I outer-membrane protein is expressed as a recombinant flagellin.

-103-

54. A method of immunizing against *Neisseria meningitidis* comprising inoculating an individual with the microorganism of claim 49.
55. A mutant meningococcal strain incapable of producing class 2 and 3 outer-membrane protein.
56. The mutant meningococcal strain H115, CBS 636.89
57. A mutant meningococcal strain, incapable of producing class 2 and 3 outer-membrane protein, transformed with a heterologous gene encoding a non-native class 1 outer-membrane protein.
58. A method of identifying vaccine epitopes of a gram negative porin protein, comprising
- obtaining the amino acid sequence of the porin protein;
 - comparing the amino acid sequence of the porin protein to the amino acid sequence of a meningococcal class 1 outer-membrane protein (OMP) to predict the location, size and sequence of exposed surface loop structures of the porin protein based upon the known corresponding structures in the class 1 OMP; and
 - identifying potential vaccine epitopes in the surface loop structures based upon the predicted location, size and sequence of the structures.

Figure 1




FIGURE 2

H4476	GLY ARG ASN TYR GLN LEU GLN LEU THR GLU ALA	20	GLN ALA ALA ASN GLY GLY ALA	30
	GCC AGG AAC TAC CAG CTG CAA TTG ACT GAA GCA		CAA GCC GCT AAC GGT GGA GCG AG	
M1080	Ile *** Ala		Gln Pro Gln Ala Thr Asn Gly Val Gln Gly	
	AT. T.. GC.C...G CAG CCC ...A A.. ...TG CAA G...			
H355	Phe		Pro Pro Ser Lys Ser Gln Pro	
	T.GC ...C.G CCC TC. AAG AG. C.A CC.			
6940	Asn Ile		Pro Pro Ser Lys Gly Gln Thr Gly	
	.AC ...ATT ...C...C.A CCC TC. AAA .GT C.G ACG ..C			
6557	Tyr		Gln Pro Ser Arg Thr Gln Gly Gln Thr	
CA. CCC TC. AGA A.. C.A ...CA. A.. ...			
870227	Ile		Pro Leu Pro Asn Ile Gln Pro Gln	
	AT.GC ...C.G CTC .C. AAT AT. C.A CC. CAG			
B40	Ile		Pro Leu Pro Asn Ile Gln Pro Gln	
	AT.GC ...C.G CTC .C. AAT AT. C.A CC. CAG			
H4476	GLY Gln Val Lys Val		Thr Lys Val Thr Lys Ala	
	GGT CAG GTA AAA GTT		ACT AAA GTT ACT AAG GCC	
M1080	Arg Gln Gly Asn Gln Val Thr		Val	
	.G. CA. GGC AA. CAG GTA .C.T.	G... ..C	
H355	Gln Val		Ala	
	CAG GTA	
6940	Asn		Ala	
	.A.	
6557	Asn Gln Val		Ala	
	AA. CAG GTA	
8700227	Arg		Arg	
	CG.	
B40	Arg		Arg	
	CG.	





SUBSTITUTE SHEET

PCT/US89/05678

41/7

SUBSTITUTE SHEET

	210	220
H4476	Lys Asn Gly Gly Phe Ala Gly Asn Tyr Ala Phe Lys Tyr Ala Arg His Ala Asn Val Gly Arg AAA AAT GGC GGT TTT GCC GGG AAC TAT GCC TTT AAA TAT GCC ACA CAC GCC AAT GTC GGA CGT	Asn
M1080	..G	Arg
H355	Lys
6940	Lys
6557	Lys
870227	Lys
B40	Lys






FIG. 4e




FIG. 4f




FIG. 4g








FIG. 4h





10/17



11/17

FIG. 5h




FIG. 5g




Figure 6. HYBRID N. MENINGIDITIS FLAGELLIN CONSTRUCTIONS






Figure 7. HYBRID N. MENINGITIDITIS FLAGELLIN STRUCTURES



13/17

FIG. 8. A representative HPLC of purified PCB12-10-6 showing a single major peak




14/17

15/17

FIG.9

A representative SDS-page of PCB12-10-6.




Lane#
1
2
3
4
5
6
7
8
9
10

Sample
Blank
High Molecular Weight Standard
Dialyzed PCB12-10-6, step 2
Fractions off peak PCB12-10-6
Purified PCB12-10-6 from HPLC, dialyzed
in PBS, 20 µg
Purified PCB12-10-6 from HPLC, dialyzed
in PBS, 1 µg
LPS - 5828
Blank
High Molecular Weight Standard
Blank

16/17

FIG.10

Photographs of representative western blot analysis of CB1 and CB2 CRM197 conjugates.



Lane#
1
2
3
4
5

Sample
Molecular Weight Standard
M20-CRM, 10 μ g
M20-CRM, 20 μ g
Molecular Weight Standard
M21-CRM, 1 μ g

SUBSTITUTE SHEET

PUTATIVE CONFORMATION MENINGOCOCCAL CLASS I OUTER-MEMBRANE PROTEIN PL.16

SUBSTITUTE SHEET




FIG. 11